

# BOUSFIELD LATTICES OF NON-NOETHERIAN RINGS: SOME QUOTIENTS AND PRODUCTS

F. LUKE WOLCOTT

**ABSTRACT.** In the context of a well generated tensor triangulated category, Section 3 investigates the relationship between the Bousfield lattice of a quotient and quotients of the Bousfield lattice. In Section 4 we develop a general framework to study the Bousfield lattice of the derived category of a commutative or graded-commutative ring, using derived functors induced by extension of scalars. Section 5 applies this work to extend results of Dwyer and Palmieri [DP08] to new non-Noetherian rings.

## 1. INTRODUCTION

Let  $R$  be a commutative ring and consider the unbounded derived category  $D(R)$  of right  $R$ -modules. Given an object  $X \in D(R)$ , define the Bousfield class  $\langle X \rangle$  of  $X$  to be  $\{W \in D(R) \mid W \otimes_R^L X = 0\}$ . Order Bousfield classes by reverse inclusion, so  $\langle 0 \rangle$  is the minimum and  $\langle R \rangle$  is the maximum. It is known that there is a set of such Bousfield classes. The join of any set  $\{\langle X_\alpha \rangle\}$  is the class  $\langle \coprod_\alpha X_\alpha \rangle$ , and the meet of a set of classes is the join of all the lower bounds. The collection of Bousfield classes thus forms a lattice, called the Bousfield lattice  $\text{BL}(D(R))$ .

A full subcategory of  $D(R)$  is localizing if it is closed under triangles and arbitrary coproducts. Thus every Bousfield class is a localizing subcategory. A result of Neeman's [Nee92] shows that when  $R$  is Noetherian, every localizing subcategory is a Bousfield lattice, and this lattice is isomorphic to the lattice of subsets of the prime spectrum  $\text{Spec } R$ .

The case of a non-Noetherian ring is much less understood. Given a ring  $k$ , fix  $n_i > 1$  and define

$$\Lambda_k = \frac{k[x_1, x_2, x_3, \dots]}{(x_1^{n_1}, x_2^{n_2}, x_3^{n_3}, \dots)},$$

and give  $\Lambda_k$  a grading by setting  $\deg(x_i) = 2^i$ . Consider the unbounded derived category  $D(\Lambda_k)$  of right graded  $\Lambda_k$ -modules; objects in  $D(\Lambda_k)$  are bi-graded. Dwyer and Palmieri [DP08] studied the Bousfield lattice of this category, when  $k$  is a countable field (see Example 2.16 below for more details). The initial motivation for the present work was to extend their main results to the case where  $k = \mathbb{Z}_{(p)}$ . We have done this fairly completely, and in the process developed tools that apply in much more general settings.

Our most general results apply to the Bousfield lattice of any well generated tensor triangulated category, and appear in Section 3. Iyengar and Krause [IK11] recently showed that a well generated tensor triangulated category has a set of

---

*Date:* January 21, 2013.

*Key words and phrases.* Bousfield lattice, non-Noetherian, derived category.

Bousfield classes, and thus a Bousfield lattice. Note that compactly generated tensor triangulated categories are well generated, and in particular those generated by the tensor unit are. For simplicity in this introduction, suppose  $\mathcal{T}$  is a tensor triangulated category generated by the tensor unit  $\mathbf{1}$ ; this includes the case of the derived category of a ring, but also the stable homotopy category and the stable module category of a  $p$ -group. Let  $- \wedge -$  denote the tensor product, and  $- \vee -$  denote the join.

The results of Section 3 concern the relationship between the quotient of a lattice and the lattice of a quotient. Given  $Z \in \mathcal{T}$ , consider the Verdier quotient  $\mathcal{T}/\langle Z \rangle$ ; this is well generated because  $\langle Z \rangle$  is. The quotient functor  $\pi : \mathcal{T} \rightarrow \mathcal{T}/\langle Z \rangle$  induces a well-defined, order-preserving map of lattices  $\pi : \mathrm{BL}(\mathcal{T}) \rightarrow \mathrm{BL}(\mathcal{T}/\langle Z \rangle)$ , where  $\langle X \rangle \mapsto \langle \pi X \rangle$ .

Given  $Z \in \mathcal{T}$ , define  $a\langle Z \rangle$  to be the join of all classes  $\langle Y \rangle$  such that  $\langle Z \wedge Y \rangle = \langle 0 \rangle$ . For any class  $\langle X \rangle \in \mathrm{BL}(\mathcal{T})$ , define  $\langle X \rangle \downarrow$  to be the collection of classes less than or equal to  $\langle X \rangle$ . In Definition 2.19 we give a notion of quotient lattice. Our first results, then, are the following.

**Proposition 3.2.** *Let  $\langle Z \rangle$  be any Bousfield class in  $\mathrm{BL}_{\mathcal{T}}$ . Then  $\pi$  induces an epimorphism of lattices with trivial kernel*

$$\bar{\pi} : \mathrm{BL}(\mathcal{T})/(\langle a\langle Z \rangle \rangle \downarrow) \longrightarrow \mathrm{BL}(\mathcal{T}/\langle Z \rangle).$$

In this context, we say a class  $\langle X \rangle$  is complemented if  $\langle X \rangle \vee a\langle X \rangle = \langle \mathbf{1} \rangle$ . (In Section 2 we discuss complemented classes when  $\mathbf{1} \notin \mathcal{T}$ .) The sub-poset of complemented classes is denoted  $\mathrm{BA}(\mathcal{T})$ .

**Corollary 3.3.** *If  $\langle Z \rangle$  is complemented then the above map is an isomorphism of lattices.*

We also consider the sub-poset  $\mathrm{DL}(\mathcal{T})$  of classes  $\langle X \rangle$  such that  $\langle X \wedge X \rangle = \langle X \rangle$ .

**Proposition 3.5.** *If  $\langle Z \rangle$  is an element of  $\mathrm{DL}(\mathcal{T})$  but is not complemented, then the map in Proposition 3.2 is not an isomorphism. This happens in the stable homotopy category and in  $D(\Lambda_k)$ , where  $k$  is a countable field.*

These results rely in part on an interesting observation that we have been unable to find in the literature. Call an object  $X \in \mathcal{T}$  square-free if  $X$  is nonzero but  $X \wedge X = 0$ .

**Corollary 2.8.** *There are no square-zero objects in  $\mathcal{T}$  if and only if*

$$\mathrm{BL} = \mathrm{DL} = \mathrm{BA}.$$

Section 4 specializes to look at functors between derived categories of rings. A ring map  $f : R \rightarrow S$  induces a functor  $f_* : \mathrm{Mod}\text{-}R \rightarrow \mathrm{Mod}\text{-}S$ , via extension of scalars, and the forgetful functor  $f^*$  is a right adjoint. This carries to the level of chain complexes, and we get an adjoint pair of derived functors on derived categories

$$f_{\bullet} : D(R) \rightleftarrows D(S) : f^{\bullet}.$$

These functors induce maps between lattices, where  $f_{\bullet}\langle X \rangle = \langle f_{\bullet}X \rangle$  and  $f^{\bullet}\langle Y \rangle = \langle f^{\bullet}Y \rangle$ , which preserve order and arbitrary joins.

First we investigate the behavior of the sub-posets  $\mathbf{BA}$  and  $\mathbf{DL}$  under  $f_\bullet$  and  $f^\bullet$ . Let  $\langle M_f \rangle$  be the join of all classes  $\langle Y \rangle$  with  $f_\bullet \langle Y \rangle = \langle 0 \rangle$ . Abbreviate  $\mathbf{BL}(D(R))$  to  $\mathbf{BL}(R)$  or  $\mathbf{BL}_R$ , and likewise for  $\mathbf{BA}$  and  $\mathbf{DL}$ . Our most general statement is the following.

**Proposition 4.15.** *Suppose  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ . The following hold.*

- (1) *The map  $f_\bullet$  sends  $\mathbf{DL}_R$  onto  $\mathbf{DL}_S$ , and the map  $f^\bullet$  injects  $\mathbf{DL}_S$  into  $\mathbf{DL}_R$ .*
- (2) *The map  $f_\bullet$  sends  $\mathbf{BA}_R$  onto  $\mathbf{BA}_S$ , and if  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$  then  $f^\bullet$  injects  $\mathbf{BA}_S$  into  $\mathbf{BA}_R$ .*

Next we establish maps between various quotients and lattices.

$$\begin{array}{ccc}
 \mathbf{BL}(R) & \xrightarrow{f_\bullet} & \mathbf{BL}(S) \\
 \downarrow & \nearrow \exists & \uparrow (\dagger) \\
 \mathbf{BL}(R)/\langle M_f \rangle \downarrow & \xrightarrow{(*)} & \mathbf{BL}(D(R)/\langle f^\bullet S \rangle)
 \end{array}$$

We show  $\langle M_f \rangle = a\langle f^\bullet S \rangle$ , and so Corollary 3.3 implies that the map  $(*)$  is an isomorphism when  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$ . Theorem 4.18 states that the map  $(\dagger)$  exists and is an isomorphism when  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ .

Finally, Section 5 applies the results of the previous two sections. Let  $g : \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{F}_p}$  be the obvious projection, and let  $h : \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{Q}}$  be inclusion. These maps give derived functors and lattice maps, as above.

The map  $g_\bullet$  has  $\langle g_\bullet g^\bullet X \rangle = \langle X \rangle$  for all  $\langle X \rangle$  (Proposition 5.3), but we show that  $h_\bullet$  does not (Lemma 5.20). The diagram above becomes the following.

$$\begin{array}{ccc}
 \mathbf{BL}(\Lambda_{\mathbb{Z}_{(p)}}) & \xrightarrow{g_\bullet} & \mathbf{BL}(\Lambda_{\mathbb{F}_p}) \\
 \downarrow & \nearrow & \uparrow \cong \\
 \mathbf{BL}(\Lambda_{\mathbb{Z}_{(p)}})/\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle \downarrow & \xrightarrow{\cong} & \mathbf{BL}(D(\Lambda_{\mathbb{Z}_{(p)}})/\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle)
 \end{array}$$

Theorem 5.14 gives a splitting of the Bousfield lattice of  $\Lambda_{\mathbb{Z}_{(p)}}$  as the product lattice  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle \downarrow \times \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle \downarrow$ . Combining this with other results, we conclude the following.

**Corollaries 5.17 and 5.18.** *The functors  $g_\bullet$  and  $h_\bullet$  induce lattice isomorphisms*

$$\mathbf{BL}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \mathbf{BL}(\Lambda_{\mathbb{F}_p}) \times \mathbf{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})),$$

$$\mathbf{DL}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \mathbf{DL}(\Lambda_{\mathbb{F}_p}) \times \mathbf{DL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})),$$

$$\mathbf{BA}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \mathbf{BA}(\Lambda_{\mathbb{F}_p}) \times \mathbf{BA}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})),$$

$$\text{where } \langle X \rangle \mapsto (g_\bullet \langle X \rangle, \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle).$$

As immediate corollaries to this, we get that the cardinality of  $\mathbf{BL}(\Lambda_{\mathbb{Z}_{(p)}})$  is  $2^{2^{\aleph_0}}$  (Corollary 5.19) and that, unlike in  $\mathbf{BL}(\Lambda_{\mathbb{F}_p})$ , in  $\mathbf{BL}(\Lambda_{\mathbb{Z}_{(p)}})$  there is no nonzero minimum Bousfield class (Proposition 5.4).

Section 2 contains background on Bousfield lattices and gives examples. With the exception of Proposition 2.6 and its corollaries, and our treatment of complemented classes when  $\mathbf{1} \notin \mathsf{T}$ , the contents are not new. The results of the Sections 3 – 5 are new, unless cited. We are grateful to John Palmieri and Dan Christensen for many helpful conversations and suggestions.

## 2. BACKGROUND ON BOUSFIELD LATTICES

In this section we review the definition and basic properties of Bousfield classes and the Bousfield lattice, and outline some of what is known about the Bousfield lattice in several examples. Most of the following general properties of Bousfield classes were first established by Bousfield [Bou79a, Bou79b] in the context of the stable homotopy category. Further work was done in [Rav84, HPS97, HP99, IK11]. Our lattice theory reference is [Bir79]. We will work in the context of a well generated tensor triangulated category, which we now define.

**Definition 2.1.** [Kra10, §6.3] Let  $\mathsf{T}$  be a triangulated category which admits arbitrary coproducts and fix a regular cardinal  $\alpha$ . An object  $X$  in  $\mathsf{T}$  is called  *$\alpha$ -small* if every morphism  $X \rightarrow \coprod_{i \in I} Y_i$  in  $\mathsf{T}$  factors through  $\coprod_{i \in J} Y_i$  for some subset  $J \subseteq I$  with  $\text{card}(J) < \alpha$ . The triangulated category  $\mathsf{T}$  is called  *$\alpha$ -well generated* if it is perfectly generated by a set of  $\alpha$ -small objects, and  $\mathsf{T}$  is called *well generated* if it is  $\beta$ -well generated for some cardinal  $\beta$ .

A category is  $\aleph_0$ -well generated if and only if it is compactly generated. A triangulated category is *tensor triangulated* if it has a symmetric monoidal product, which we will denote  $- \wedge -$ , that is compatible with the triangulation, is exact in both variables, and commutes with arbitrary coproducts [HPS97, App. A]. We will denote the tensor unit by  $\mathbf{1}$ , and assume that  $\mathbf{1}$  is compact. We do not assume that  $\mathbf{1} \in \mathsf{T}$ .

Henceforth, let  $\mathsf{T}$  denote a well generated tensor triangulated category. Let  $\Sigma$  denote the shift.

**Definition 2.2.** Let  $W, X$ , and  $Y$  be objects in  $\mathsf{T}$ .

- (1) We say  $W$  is  *$X$ -acyclic* if  $W \wedge X = 0$ .
- (2) The collection of  $X$ -acyclics is denoted  $\langle X \rangle$  and called the *Bousfield class* of  $X$ .
- (3) We say  $X$  and  $Y$  are *Bousfield equivalent* if they have the same acyclics.

There is a partial ordering on Bousfield classes, given by reverse inclusion. So we say

$$\langle X \rangle \leq \langle Y \rangle \text{ if and only if } (W \wedge Y = 0 \implies W \wedge X = 0).$$

Note that  $\langle 0 \rangle$  is the minimum class under this ordering, and  $\langle X \rangle = \langle 0 \rangle$  if and only if  $X = 0$ . The join of a set of classes  $\{\langle X_\alpha \rangle\}_{\alpha \in A}$  is given by

$$\bigvee_{\alpha \in A} \langle X_\alpha \rangle := \left\langle \prod_{\alpha \in A} X_\alpha \right\rangle.$$

It was recently shown in [IK11, Thm. 3.1] that in a well generated tensor triangulated category there is always a set of Bousfield classes. We can define the meet

(denoted  $\wedge$ ) of any set of classes  $\{\langle X_\alpha \rangle\}$  to be the join of all the lower bounds; this join is over a set, and a nonempty set because  $\langle 0 \rangle$  is the minimum.

A partially ordered set with finite joins and meets is called a *lattice*. A lattice with arbitrary joins and meets is *complete*. The collection of Bousfield classes of  $\mathbf{T}$  is thus a complete lattice, called the *Bousfield lattice*, and denoted  $\mathbf{BL}$ .

In any complete lattice there is also a maximum element  $\langle \mathbf{Max} \rangle$ , given by joining all elements. When  $\mathbf{1} \in \mathbf{T}$ , then clearly  $\langle \mathbf{Max} \rangle = \langle \mathbf{1} \rangle = \{0\}$ . This is the case in the derived category  $D(R)$  of a ring, of course, since  $R \in D(R)$ . Most discussions of Bousfield lattices assume  $\mathbf{1} \in \mathbf{T}$ , but we work slightly more generally in this and the following section, and at the end of the paper. The reason for this is that we would like to consider Bousfield lattices of certain subcategories  $\mathbf{S} \subseteq \mathbf{T}$ , where  $\mathbf{1} \notin \mathbf{S}$ . For example, see Remark 2.12 and Corollary 5.17.

The tensor product gives another operation on Bousfield classes,

$$\langle X \rangle \wedge \langle Y \rangle := \langle X \wedge Y \rangle.$$

We always have  $\langle X \rangle \wedge \langle Y \rangle \leq \langle X \rangle \wedge \langle Y \rangle$ .

**Definition 2.3.**

- (1) Define  $\mathbf{DL} = \{\langle X \rangle \in \mathbf{BL} \text{ with } \langle X \rangle = \langle X \wedge X \rangle\}$ .
- (2) A Bousfield class  $\langle X \rangle$  is called *complemented* if there exists a class  $\langle X^c \rangle$  such that  $\langle X \rangle \wedge \langle X^c \rangle = \langle 0 \rangle$  and  $\langle X \rangle \vee \langle X^c \rangle = \langle \mathbf{Max} \rangle$ . Call  $\langle X^c \rangle$  a *complement* of  $\langle X \rangle$ .
- (3) Define  $\mathbf{BA}$  to be the collection of Bousfield classes in  $\mathbf{DL}$  that are complemented and have a complement in  $\mathbf{DL}$ .

When the category needs to be specified, we will write  $\mathbf{BL}_{\mathbf{T}}$ ,  $\mathbf{DL}_{\mathbf{T}}$ , and  $\mathbf{BA}_{\mathbf{T}}$ , or  $\mathbf{BL}(\mathbf{T})$ , etc. In the case where  $\mathbf{T} = D(R)$  is the derived category of a ring, we will use the notation  $\mathbf{BL}_R$ ,  $\mathbf{DL}_R$ , and  $\mathbf{BA}_R$ , or  $\mathbf{BL}(R)$ , etc. instead.

The sub-poset  $\mathbf{DL} \subseteq \mathbf{BL}$  is closed under arbitrary joins, and under the tensor operation, but not under meets; the meet in  $\mathbf{BL}$  of two elements of  $\mathbf{DL}$  may not be in  $\mathbf{DL}$ . However, when we restrict to  $\mathbf{DL}$ , the meet is given by tensoring: if  $\langle X \rangle, \langle Y \rangle, \langle Z \rangle \in \mathbf{DL}$  have  $\langle Z \rangle \leq \langle X \rangle$  and  $\langle Z \rangle \leq \langle Y \rangle$ , then  $\langle Z \rangle = \langle Z \wedge Z \rangle \leq \langle X \wedge Y \rangle$ , so  $\langle X \wedge Y \rangle$  is the greatest lower bound. A lattice is called *distributive* if meets distribute across finite joins (equiv. if joins distribute across finite meets; see [Bir79, I.6, Thm. 9]); it is a *frame* if meets distribute across arbitrary joins. Since the tensor product commutes with arbitrary coproducts,  $\mathbf{DL}$  is a frame.

In general, a complemented class may have multiple complements. When  $\mathbf{1} \in \mathbf{T}$ , every complemented class is in  $\mathbf{DL}$ , because then  $\langle \mathbf{Max} \rangle = \langle \mathbf{1} \rangle$  and we have

$$\langle X \rangle = \langle X \wedge \mathbf{1} \rangle = \langle X \rangle \wedge (\langle X \rangle \vee \langle X^c \rangle) = (\langle X \rangle \wedge \langle X \rangle) \vee (\langle X \rangle \wedge \langle X^c \rangle) = \langle X \wedge X \rangle.$$

Furthermore, if  $\langle X \rangle \in \mathbf{BA}$  then  $\langle X \rangle$  has a unique complement in  $\mathbf{DL}$ . Indeed, if  $\langle X^c \rangle, \langle \widetilde{X}^c \rangle \in \mathbf{DL}$  are two complements, then since the meet is given by tensoring, we have

$$\langle X^c \rangle = \langle X^c \rangle \wedge (\langle X \rangle \vee \langle X^c \rangle) = \langle X^c \rangle \wedge (\langle X \rangle \vee \langle \widetilde{X}^c \rangle) = \langle X^c \rangle \wedge \langle \widetilde{X}^c \rangle,$$

and likewise  $\langle \widetilde{X}^c \rangle = \langle X^c \rangle \wedge \langle \widetilde{X}^c \rangle$ .

One can check that  $\mathbf{BA}$  is a sublattice of  $\mathbf{DL}$  (i.e. is closed under finite joins and meets), with  $\langle (X \vee Y)^c \rangle = \langle X^c \rangle \wedge \langle Y^c \rangle$  and  $\langle (X \wedge Y)^c \rangle = \langle X^c \rangle \vee \langle Y^c \rangle$ . In general, however,  $\mathbf{BA}$  is not closed under infinite joins. A *Boolean algebra* is a distributive

lattice in which every element is complemented; thus  $\mathbf{BA}$  is a Boolean algebra, and this explains the notation.

We can use the tensor product to define another operation on Bousfield classes.

**Definition 2.4.** For any Bousfield class  $\langle Z \rangle$  in the Bousfield lattice  $\mathbf{BL}$ , define the *complementation operator*  $a(-)$  to be

$$a\langle Z \rangle = \bigvee_{\langle Y \wedge Z \rangle = \langle 0 \rangle} \langle Y \rangle.$$

The complementation operator was first considered by Bousfield [Bou79a], and later by Hovey and Palmieri [HP99], in the stable homotopy category. Note that the definition requires knowing there is a set of Bousfield classes. They prove the following properties of  $a(-)$  in that context, but the proof is formal and applies in any well generated tensor triangulated category.

**Lemma 2.5.** [HP99, Lemma 2.3] *The complementation operator  $a(-)$  has the following properties.*

- (1)  $\langle E \rangle \leq a\langle X \rangle$  if and only if  $\langle E \rangle \wedge \langle X \rangle = \langle 0 \rangle$ .
- (2)  $a(-)$  is order-reversing:  $\langle X \rangle \leq \langle Y \rangle$  if and only if  $a\langle X \rangle \geq a\langle Y \rangle$ .
- (3)  $a^2\langle X \rangle = \langle X \rangle$ .

Note that we always have  $\langle X \rangle \wedge a\langle X \rangle = \langle 0 \rangle$  and  $\langle X \rangle \wedge a\langle X \rangle \leq \langle \mathbf{Max} \rangle$ . If  $\langle X \rangle \in \mathbf{DL}$ , then  $a\langle X \rangle$  is not necessarily in  $\mathbf{DL}$ . If  $\langle X \rangle$  is complemented, with some complement  $\langle X^c \rangle$ , then  $\langle X \rangle$  is also complemented by  $a\langle X \rangle$ . This is because, by the Lemma,  $\langle X \rangle \wedge \langle X^c \rangle = \langle 0 \rangle$  implies  $\langle X^c \rangle \leq a\langle X \rangle$ , and thus  $\langle \mathbf{Max} \rangle = \langle X \rangle \vee \langle X^c \rangle \leq \langle X \rangle \vee a\langle X \rangle$ . It follows that if  $\mathbf{1} \in \mathbf{T}$  and  $\langle X \rangle \in \mathbf{BA}$ , then  $a\langle X \rangle$  is in  $\mathbf{DL}$  and is the unique complement of  $\langle X \rangle$ .

We briefly mention a surprising but simple result using complementation, which we have been unable to find in the literature. Call an object  $X \in \mathbf{T}$  *square-zero* if  $X$  is nonzero but  $X \wedge X = 0$ .

**Proposition 2.6.** *If there are no square-zero objects in  $\mathbf{T}$ , then every object is complemented.*

**Proof.** Let  $X \in \mathbf{T}$  be arbitrary. It suffices to show that  $\langle X \rangle \vee a\langle X \rangle \geq \langle \mathbf{Max} \rangle$ . Suppose  $Y$  has  $\langle Y \rangle \wedge \langle X \rangle = \langle 0 \rangle$  and  $\langle Y \rangle \wedge a\langle X \rangle = \langle 0 \rangle$ . Part (1) of Lemma 2.5 implies that  $\langle Y \rangle \leq \langle X \rangle$ , and from this we conclude that  $Y \wedge Y = 0$ . Our assumption forces  $Y = 0$  so  $\langle Y \rangle \wedge \langle \mathbf{Max} \rangle = \langle 0 \rangle$ . Thus  $\langle X \rangle$  is complemented by  $a\langle X \rangle$ .  $\square$

**Corollary 2.7.** *If  $\mathbf{DL} = \mathbf{BL}$ , then  $\mathbf{BA} = \mathbf{DL} = \mathbf{BL}$ .*

**Corollary 2.8.** *If  $\mathbf{1} \in \mathbf{T}$ , then there are no square-zero objects in  $\mathbf{T}$  if and only if  $\mathbf{BA} = \mathbf{DL} = \mathbf{BL}$ .*

**2.1. Subcategories and quotient categories.** Well generated categories behave well under taking subcategories and quotients.

**Definition 2.9.** Let  $X$  be an object in  $\mathbf{T}$ .

- (1) A full subcategory  $\mathbf{S} \subseteq \mathbf{T}$  is *thick* if it is closed under triangles and retracts.
- (2) The smallest thick subcategory containing  $X$  is denoted  $\mathbf{th}(X)$ ; this is also called the *thick subcategory generated by  $X$* .

- (3) A full subcategory  $\mathcal{S} \subseteq \mathcal{T}$  is *localizing* if it is closed under triangles, retracts, and arbitrary coproducts.
- (4) The smallest localizing subcategory containing  $X$  is denoted  $\text{loc}(X)$ ; this is also called the *localizing subcategory generated by  $X$* .

Note that every Bousfield class is a localizing subcategory. A localizing subcategory  $\mathcal{S} \subseteq \mathcal{T}$  is well generated if and only if  $\mathcal{S} = \text{loc}(X)$  for some  $X \in \mathcal{T}$  [IK11, Rmk. 2.2].

**Lemma 2.10.** *Every Bousfield class  $\langle Z \rangle \subseteq \mathcal{T}$  is well generated. Thus for all  $Z \in \mathcal{T}$ , there exists an element  $aZ \in \mathcal{T}$  such that  $\langle Z \rangle = \text{loc}(aZ)$ .*

**Proof.** This follows from Proposition 2.1 in [IK11], since  $\langle Z \rangle$  is the kernel of the exact coproduct-preserving functor  $F = (- \wedge Z) : \mathcal{T} \rightarrow \mathcal{T}$ .  $\square$

**Lemma 2.11.** *For any  $Z \in \mathcal{T}$ , we have  $\langle aZ \rangle = a\langle Z \rangle$ .*

**Proof.** Because  $aZ \in \langle Z \rangle$ , Lemma 2.5 implies that  $\langle aZ \rangle \leq a\langle Z \rangle$ . If  $\langle Y \rangle$  has  $\langle Y \wedge Z \rangle = \langle 0 \rangle$ , then  $Y \in \langle Z \rangle = \text{loc}(aZ)$ . It follows that  $\langle Y \rangle \leq \langle aZ \rangle$ . Therefore  $a\langle Z \rangle \leq \langle aZ \rangle$  and equality holds.  $\square$

Given any well generated localizing subcategory  $\mathcal{S} \subseteq \mathcal{T}$ , we can consider the Bousfield lattice  $\text{BL}(\mathcal{S})$ . Some care is necessary, since for  $X \in \mathcal{S}$ , the Bousfield class  $\langle X \rangle$  in  $\text{BL}(\mathcal{S})$  is  $\{W \in \mathcal{S} \mid X \wedge W = 0\}$ . If  $X, Y \in \mathcal{S}$  have  $\langle X \rangle \leq \langle Y \rangle$  in  $\text{BL}(\mathcal{S})$ , it does not necessarily follow that  $\langle X \rangle \leq \langle Y \rangle$  in  $\text{BL}(\mathcal{T})$ . However, see Lemmas 5.11 and 5.12.

**Remark 2.12.** If  $\mathcal{S} \subseteq \mathcal{T}$  is a well generated localizing subcategory, then  $\mathcal{S} = \text{loc}(X)$  for some  $X \in \mathcal{T}$ , and in this case  $\langle \text{Max} \rangle$  in  $\text{BL}(\mathcal{S})$  is  $\langle X \rangle$ . This is because  $Y \in \text{loc}(X)$  always implies  $\langle Y \rangle \leq \langle X \rangle$ .

If  $\mathcal{S} \subseteq \mathcal{T}$  is any localizing subcategory, we can form the Verdier quotient  $\mathcal{T}/\mathcal{S}$ . This category has a tensor triangulated structure induced by that on  $\mathcal{T}$ , such that the quotient functor  $\pi : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{S}$  is exact, and  $\pi(\mathbf{1}_{\mathcal{T}}) = \mathbf{1}_{\mathcal{T}/\mathcal{S}}$ . If  $\mathcal{S}$  is well generated, then so is  $\mathcal{T}/\mathcal{S}$ , by [Nee01, Cor. 4.4.3] or [Kra10, Thm. 7.2.1]. This also implies that  $\mathcal{T}/\mathcal{S}$  has Hom sets.

**2.2. Examples.** Next we survey several examples of categories and their Bousfield lattices.

**Example 2.13.** Let  $R$  be a commutative ring, or a graded-commutative ring. Let  $D(R)$  denote the unbounded derived category of right  $R$ -modules, or of right graded  $R$ -modules. If  $R$  is graded, we think of objects in  $D(R)$  as bi-graded; in either case we assume the differential decreases the chain degree by one. Then  $D(R)$  is a tensor triangulated category, with the product  $A \wedge B = A \otimes_R^L B$  given by the left derived tensor product [HPS97, §9.3]. The tensor unit is the module  $R$  thought of as a complex concentrated in degree zero. Furthermore,  $D(R) = \text{loc}(R)$ , so  $D(R)$  is compactly generated, hence well generated. The Bousfield lattice of  $D(R)$  is well-understood when  $R$  is Noetherian; see the next example. When  $R$  is non-Noetherian our understanding of the Bousfield lattice is limited to several specific rings; see Example 2.16.

**Example 2.14.** Iyengar and Krause [IK11] investigate the Bousfield lattice of a compactly generated tensor triangulated category that is stratified by the action

of a graded Noetherian ring  $R$ . This general setting, developed in [BIK08, BIK11], building on [Nee92, BCR97, HPS97], includes the unbounded derived category of a commutative Noetherian ring; the stable module category  $\text{StMod}(kG)$  of a finite group, where the characteristic of  $k$  divides the order of the group, and then also the homotopy category  $K(\text{Inj } kG)$  of complexes of injectives; and DG modules over a formal commutative DG algebra with a Noetherian cohomology ring. They show that in such a category the Bousfield lattice is isomorphic to the lattice of subsets of the homogeneous prime spectrum of  $R$ , and  $\text{BA} = \text{DL} = \text{BL}$ . In the case of a commutative Noetherian ring  $R$ ,  $D(R)$  is stratified by  $R$ , and so  $\text{BL}_R$  is isomorphic to the lattice of subsets of  $\text{Spec } R$ . The isomorphism is given in terms of support.

**Example 2.15.** The  $(p\text{-local})$  stable homotopy category  $\mathcal{S}$  is a tensor triangulated category, with the product the smash product, and the unit the  $(p\text{-local})$  sphere spectrum  $S^0$ . Since  $\mathcal{S} = \text{loc}(S^0)$ , this category is well generated. Bousfield [Bou79a] showed that the class of every finite spectrum is in  $\text{BA}$ , the class of every ring spectrum is in  $\text{DL}$ , but for example the class of  $H\mathbb{Z}$  is in  $\text{DL}$  but not in  $\text{BA}$ . He also showed that the Brown-Comenetz dual  $IS^0$  of the sphere has  $IS^0 \wedge IS^0 = 0$ , so  $\text{DL} \subsetneq \text{BL}$ . Hovey and Palmieri [HP99] study finer structure of the Bousfield lattice of this category.

**Example 2.16.** Fix a countable field  $k$  and integers  $n_i > 1$ , and consider the ring

$$\Lambda := \frac{k[x_1, x_2, \dots]}{(x_1^{n_1}, x_2^{n_2}, \dots)},$$

with the  $x_i$  graded so that  $\Lambda$  is graded-connected and finite-dimensional in each degree. Let  $D(\Lambda)$  be the derived category of graded  $\Lambda$ -modules; objects in  $D(\Lambda)$  are bigraded. Neeman [Nee00] first considered such a ring (with  $n_i = i$ ), showing the Bousfield lattice is large, although the homogeneous prime spectrum is trivial. Dwyer and Palmieri [DP08] examine the Bousfield lattice of  $D(\Lambda)$  in depth. They show the Bousfield lattice has cardinality exactly  $2^{2^{\aleph_0}}$ .

Let  $I\Lambda = \text{Hom}_k^*(\Lambda, k)$  be the graded vector-space dual of  $\Lambda$ . This is a  $\Lambda$ -module, and we consider it as an object of  $D(\Lambda)$  concentrated at chain degree zero. The module  $I\Lambda$  plays an important role in [DP08]. One computation gives  $I\Lambda \wedge I\Lambda = 0$ , so  $\text{DL}_\Lambda \subsetneq \text{BL}_\Lambda$ . This is relevant, because it implies that there is no Noetherian ring that stratifies  $D(\Lambda)$ .

Furthermore,  $\langle I\Lambda \rangle$  is a minimum nonzero Bousfield class: Corollary 7.3 in [DP08] shows that for any non-zero  $E$  in  $D(\Lambda)$ , we have that  $\langle I\Lambda \rangle \leq \langle E \rangle$ . This implies that  $\text{BA}_\Lambda$  is trivial, i.e. the only complemented pair is  $\langle 0 \rangle$  and  $\langle \Lambda \rangle = \langle \mathbf{1} \rangle$  (see Prop. 5.4).

We mention one more difference among the Bousfield lattices in these examples. One can easily check that every Bousfield class is a localizing subcategory. Hovey and Palmieri [HP99, Conj. 9.1] conjecture that the converse holds in the stable homotopy category, but no progress has been made on this question. In a category that is stratified by the action of a Noetherian ring, it is indeed the case that every localizing subcategory is a Bousfield class [IK11, Cor. 4.5]. On the other hand, Greg Stevenson [Ste12], working in the unbounded derived category of a non-Noetherian ring (specifically any absolutely flat ring which is not semi-artinian), recently exhibited a localizing subcategory that is not a Bousfield class.

**2.3. Some (more) lattice theory.** Here we recall some terminology and facts from lattice theory that we will need; our reference is [Bir79]. A *sub-poset*  $K$  of a



lattice  $\mathbf{L}$  is a subset of  $\mathbf{L}$  along with the induced partial ordering. A sub-poset  $\mathbf{K}$  of a lattice  $\mathbf{L}$  is a *sublattice* if it is closed under finite joins and meets.

If  $\mathbf{K}$  and  $\mathbf{L}$  are lattices, a *lattice morphism* or *lattice map*  $F : \mathbf{K} \rightarrow \mathbf{L}$  is a set map that preserves order (so  $x \leq y$  implies  $Fx \leq Fy$ ). A *lattice epimorphism* is a lattice morphism that is onto. A *lattice isomorphism* is a lattice morphism that is a set bijection and has an order-preserving inverse.

We do not assume that a lattice morphism preserves joins or meets, or minimum and maximum elements. Nor do we assume that a lattice morphism between Bousfield lattices will commute with the tensor product operation  $\langle X \rangle \wedge \langle Y \rangle$ .

Any poset can be thought of as a category, where  $x \leq y$  if and only if there is a (unique) morphism from  $x$  to  $y$ . Joins are colimits and meets are limits. Then a complete lattice corresponds to a category that is complete and cocomplete in the categorical sense. A lattice morphism is just a functor between such categories.

**Definition 2.17.** For any element  $a$  in a lattice  $\mathbf{L}$ , define  $a\downarrow = \{x \in \mathbf{L} \mid x \leq a\}$  and  $a\uparrow = \{x \in \mathbf{L} \mid x \geq a\}$ . Note that these are both sublattices of  $\mathbf{L}$ .

**Definition 2.18.** A nonempty subset  $J$  of a lattice  $\mathbf{L}$  is an *ideal* if it is closed under finite joins, and  $a \in J$  and  $x \in \mathbf{L}$  with  $x \leq a$  implies  $x \in J$ . An ideal is *complete* if it is closed under arbitrary joins. Note that  $a\downarrow$  is an ideal, for all  $a \in \mathbf{L}$ . An ideal  $J$  is *principal* if  $J = a\downarrow$  for some  $a \in \mathbf{L}$ . Note that an ideal  $J$  is principal if and only if it is complete.

**Definition 2.19.** Given an ideal  $J$  of a lattice  $\mathbf{L}$ , and  $a, b \in \mathbf{L}$ , we say  $a \equiv b \pmod{J}$ , and write  $[a] = [b]$ , if  $a \vee c = b \vee c$  for some  $c \in J$ . The equivalence classes under this equivalence relation, with the ordering, join, and meet induced by  $\mathbf{L}$ , form a lattice  $\mathbf{L}/J$ , called the *quotient lattice*. The quotient map  $\mathbf{L} \rightarrow \mathbf{L}/J$  sending  $x \mapsto [x]$  is a lattice epimorphism.

It is not hard to show that if  $J = a\downarrow$  is a principal ideal in a lattice  $\mathbf{L}$ , then  $[x] = [y]$  in  $\mathbf{L}/J$  if and only if  $x \vee a = y \vee a$ . Every quotient of a lattice by a principal ideal is isomorphic to a sublattice: for all  $a \in \mathbf{L}$ , there is an isomorphism of lattices  $\mathbf{L}/a\downarrow \xrightarrow{\sim} a\uparrow$ , given by  $[x] \mapsto x \vee a$ .

**Definition 2.20.** Given lattices  $\mathbf{K}$  and  $\mathbf{L}$ , the product lattice is defined as the set product  $\mathbf{K} \times \mathbf{L}$ , with  $(a, b) \leq (c, d)$  precisely when  $a \leq c$  and  $b \leq d$ , and joins and meets defined termwise. One can check that, for example,  $0 \times \mathbf{L}$  is a principal ideal in  $\mathbf{K} \times \mathbf{L}$ , and there is a lattice isomorphism  $(\mathbf{K} \times \mathbf{L})/(0 \times \mathbf{L}) \cong \mathbf{K}$ .

### 3. LATTICES AND QUOTIENTS

In this section we give some results comparing the quotient of a Bousfield lattice to the Bousfield lattice of a quotient. Again, let  $\mathbf{T}$  be a well generated tensor triangulated category. Let  $Z$  be an element of  $\mathbf{T}$ , and consider the Verdier quotient  $\mathbf{T}/\langle Z \rangle$  and quotient functor  $\pi : \mathbf{T} \rightarrow \mathbf{T}/\langle Z \rangle$ .

**Lemma 3.1.** *The functor  $\pi$  induces a lattice epimorphism that preserves arbitrary joins,*

$$\pi : \mathrm{BL}(\mathbf{T}) \rightarrow \mathrm{BL}(\mathbf{T}/\langle Z \rangle), \text{ where } \langle X \rangle \mapsto \langle \pi X \rangle.$$

**Proof.** We will show that if  $X, Y \in \mathbf{T}$  have  $\langle X \rangle \leq \langle Y \rangle$  in  $\mathbf{BL}_{\mathbf{T}}$ , then  $\langle \pi X \rangle \leq \langle \pi Y \rangle$  in  $\mathbf{BL}(\mathbf{T}/\langle Z \rangle)$ . This will show that  $\pi$  is order-preserving, and by symmetry will also show that  $\pi$  is well-defined. Take  $W \in \mathbf{T}/\langle Z \rangle$  with  $W \wedge \pi Y = 0$ . Take  $\widetilde{W}$  in  $\mathbf{T}$  so  $\pi \widetilde{W} = W$ . The tensor structure on  $\mathbf{T}/\langle Z \rangle$  is such that  $\pi(\widetilde{W} \wedge Y) = \pi \widetilde{W} \wedge \pi Y = 0$ , so we have  $\widetilde{W} \wedge Y \in \langle Z \rangle$ , i.e.  $\widetilde{W} \wedge Y \wedge Z = 0$ . Then  $(\widetilde{W} \wedge Z) \wedge X = 0$ , by hypothesis, so  $\pi(\widetilde{W} \wedge X) = 0$ . This shows that  $W \wedge \pi X = 0$ .

Since  $\pi$  commutes with arbitrary coproducts, it commutes with arbitrary joins.

□

**Proposition 3.2.** *Let  $\langle Z \rangle$  be any Bousfield class in  $\mathbf{BL}_{\mathbf{T}}$ . Then  $\pi$  induces an epimorphism of lattices that preserves arbitrary joins,*

$$\bar{\pi} : \mathbf{BL}_{\mathbf{T}}/(\langle Z \rangle)_{\downarrow} \longrightarrow \mathbf{BL}(\mathbf{T}/\langle Z \rangle),$$

*such that if  $\bar{\pi}[\langle X \rangle] = \langle 0 \rangle$ , then  $[\langle X \rangle] = [\langle 0 \rangle]$ .*

**Proof.** First we show that  $\bar{\pi}$  is order-preserving. Suppose  $\langle X \rangle \leq \langle Y \rangle$  in  $\mathbf{BL}_{\mathbf{T}}/(\langle Z \rangle)_{\downarrow}$ ; this is equivalent to assuming  $\langle X \rangle \vee \langle Z \rangle \leq \langle Y \rangle \vee \langle Z \rangle$ . We want to show that  $\langle \pi X \rangle \leq \langle \pi Y \rangle$  in  $\mathbf{BL}(\mathbf{T}/\langle Z \rangle)$ . Take  $W \in \mathbf{T}/\langle Z \rangle$  with  $W \wedge \pi Y = 0$ , and let  $\widetilde{W} \in \mathbf{T}$  be such that  $\pi \widetilde{W} = W$ . Then  $0 = \pi \widetilde{W} \wedge \pi Y = \pi(\widetilde{W} \wedge Y)$ , so  $\widetilde{W} \wedge Y \in \langle Z \rangle$  and  $\widetilde{W} \wedge Y \wedge Z = \widetilde{W} \wedge Z \wedge Y = 0$ .

Since  $\langle Z \rangle \wedge \langle Z \rangle = \langle 0 \rangle$ , we also have  $\langle \widetilde{W} \wedge Z \rangle \wedge \langle Z \rangle = \langle 0 \rangle$ . Our assumption then implies that  $(\widetilde{W} \wedge Z) \wedge X = 0$ . Therefore  $\widetilde{W} \wedge X \in \langle Z \rangle$ , which says that  $0 = \pi(\widetilde{W} \wedge X) = W \wedge \pi X$ .

Thus  $\bar{\pi}$  is order-preserving, and hence well-defined. It is clearly an epimorphism.

Now suppose  $\bar{\pi}[\langle X \rangle] = \langle 0 \rangle$ . Then  $\langle \pi X \rangle = \langle 0 \rangle$ , so  $X \in \langle Z \rangle$ , i.e.  $X \wedge Z = 0$ . Using Lemma 2.5 this implies that  $\langle X \rangle \leq \langle Z \rangle$ , so  $[\langle X \rangle] = [\langle 0 \rangle]$ . □

To be a lattice isomorphism,  $\bar{\pi}$  must have an order-preserving inverse. In the remainder of this section, we will give examples of when this does and does not happen.

**Corollary 3.3.** *Let  $\langle Z \rangle$  and  $\langle Z^c \rangle$  be a pair of complemented classes in  $\mathbf{BL}_{\mathbf{T}}$ , and assume that  $\mathbf{1} \in \mathbf{T}$ . Then  $\pi$  induces a lattice isomorphism*

$$\bar{\pi} : \mathbf{BL}_{\mathbf{T}}/\langle Z^c \rangle_{\downarrow} \longrightarrow \mathbf{BL}(\mathbf{T}/\langle Z \rangle).$$

**Proof.** Recall that  $\mathbf{1} \in \mathbf{T}$  implies that complements are unique and  $\langle Z^c \rangle = a\langle Z \rangle$ . We claim that  $\psi(\langle \pi X \rangle) = [\langle X \rangle]$  is a well-defined, order-preserving inverse to  $\bar{\pi}$ . Thus we wish to show that if  $\langle \pi X \rangle \leq \langle \pi Y \rangle$ , then  $\langle X \rangle \vee \langle Z^c \rangle \leq \langle Y \rangle \vee \langle Z^c \rangle$ . By symmetry, this will also show that  $\psi$  is well-defined; by inspection, then, it is an inverse to  $\bar{\pi}$ .

Take  $W \in \mathbf{T}$  with  $W \in \langle Y \rangle \vee \langle Z^c \rangle$ . Then  $W \wedge Y = 0$ , so  $W \wedge Y \wedge Z = 0$ , i.e.  $(W \wedge Y) \in \langle Z \rangle$ . This says that  $\pi(W \wedge Y) = 0$  in  $\mathbf{T}/\langle Z \rangle$ , so  $\pi W \wedge \pi Y = 0$ . By hypothesis, this means  $\pi W \wedge \pi X = 0$ , which working backwards implies that  $W \wedge X \wedge Z = 0$ .

On the other hand, we also know that  $W \wedge Z^c = 0$ , so  $W \wedge X \wedge Z^c = 0$ . Therefore  $(W \wedge X) \in \langle Z \rangle \vee \langle Z^c \rangle = \langle \mathbf{Max} \rangle = \langle \mathbf{1} \rangle$ , so  $W \wedge X = 0$ . Thus  $W \in \langle X \vee Z^c \rangle$  as desired. □

For example, if  $L : \mathbf{T} \rightarrow \mathbf{T}$  is a smashing localization functor with colocalization  $C$ , and  $\mathbf{1} \in \mathbf{T}$ , then  $\langle L\mathbf{1} \rangle$  and  $\langle C\mathbf{1} \rangle$  are a complemented pair. This result relates

the Bousfield lattice of  $\mathcal{T}$  to the Bousfield lattice of the  $L$ -local category, which is equivalent to  $\mathcal{T}/\langle L\mathbf{1} \rangle$ . See also [IK11, Prop. 6.12].

**Corollary 3.4.** *Suppose  $\mathbf{BL}_{\mathcal{T}} = \mathbf{DL}_{\mathcal{T}}$ . Then for every Bousfield class  $\langle Z \rangle$ , the functor  $\pi$  induces a lattice isomorphism*

$$\bar{\pi} : \mathbf{BL}_{\mathcal{T}}/(\langle a\langle Z \rangle \rangle) \longrightarrow \mathbf{BL}(\mathcal{T}/\langle Z \rangle).$$

**Proof.** This follows immediately from Corollaries 2.7 and 3.3 if  $\mathbf{1} \in \mathcal{T}$ , but we will prove it more generally. As in the last proof, we claim that  $\psi(\langle \pi X \rangle) = [\langle X \rangle]$  is a well-defined, order-preserving inverse to  $\bar{\pi}$ . Suppose  $\langle \pi X \rangle \leq \langle \pi Y \rangle$ ; it suffices to show that  $\langle X \rangle \vee a\langle Z \rangle \leq \langle Y \rangle \vee a\langle Z \rangle$ .

Take  $W \in \mathcal{T}$  with  $W \in \langle Y \rangle \vee a\langle Z \rangle$ . As in the last proof,  $W \wedge Y = 0$  implies  $W \wedge X \wedge Z = 0$ . Then  $\langle W \rangle \wedge a\langle Z \rangle = \langle 0 \rangle$  implies  $\langle W \rangle \leq \langle Z \rangle$ , by Lemma 2.5. Therefore  $\langle W \wedge X \rangle \in \langle W \rangle$ . Since  $\mathbf{BL}_{\mathcal{T}} = \mathbf{DL}_{\mathcal{T}}$ , we have  $X \in \langle W \wedge W \rangle = \langle W \rangle$ , so  $W \wedge X = 0$  and this concludes the proof.  $\square$

The last two corollaries apply when  $\mathcal{T}$  is a stratified category, as discussed in Example 2.14. The next result, however, shows that  $\bar{\pi}$  fails to be an isomorphism in both the stable homotopy category and the category  $D(\Lambda)$ .

**Proposition 3.5.** *Assume  $\mathbf{1} \in \mathcal{T}$ . Suppose  $\langle Z \rangle$  in  $\mathbf{BL}_{\mathcal{T}}$  is an element of  $\mathbf{DL}_{\mathcal{T}} \setminus \mathbf{BA}_{\mathcal{T}}$ . Then the epimorphism induced by  $\pi$  in Proposition 3.2 is not an isomorphism. This happens in both the stable homotopy category and  $D(\Lambda)$ .*

**Proof.** Since  $\langle Z \rangle \notin \mathbf{BA}_{\mathcal{T}}$ , we know  $\langle Z \rangle \vee a\langle Z \rangle < \langle \mathbf{1} \rangle$ . We will show that  $\langle \pi Z \rangle = \langle \pi \mathbf{1} \rangle$  in  $\mathbf{BL}(\mathcal{T}/\langle Z \rangle)$ , but  $[\langle Z \rangle] \neq [\langle \mathbf{1} \rangle]$  in  $\mathbf{BL}_{\mathcal{T}}/(\langle a\langle Z \rangle \rangle)$ .

Since  $\pi$  is order-preserving, we know that  $\langle \pi Z \rangle \leq \langle \pi \mathbf{1} \rangle$ . We must show  $\langle \pi Z \rangle \geq \langle \pi \mathbf{1} \rangle$ . Suppose  $W \in \mathcal{T}/\langle Z \rangle$  has  $W \wedge \pi Z = 0$ . Choose  $\widetilde{W} \in \mathcal{T}$  such that  $\pi \widetilde{W} = W$ . Then  $\pi(\widetilde{W} \wedge Z) = 0$ , so  $\widetilde{W} \wedge Z \wedge Z = 0$ . This says  $\widetilde{W} \in \langle Z \wedge Z \rangle$ , and by hypothesis  $\langle Z \wedge Z \rangle = \langle Z \rangle$ , so  $\widetilde{W} \wedge Z = 0$ . Therefore  $\widetilde{W} = \widetilde{W} \wedge \mathbf{1} \in \langle Z \rangle$  and  $\pi(\widetilde{W} \wedge \mathbf{1}) = W \wedge \pi \mathbf{1} = 0$  in  $\mathcal{T}/\langle Z \rangle$ . This shows  $\langle \pi \mathbf{1} \rangle = \langle \pi Z \rangle$ .

By assumption,  $\langle Z \rangle$  is such that  $\langle Z \rangle \vee a\langle Z \rangle < \langle \mathbf{1} \rangle$ . But  $\langle \mathbf{1} \rangle = \langle \mathbf{1} \rangle \vee a\langle Z \rangle$ , so  $\langle Z \rangle \vee a\langle Z \rangle < \langle \mathbf{1} \rangle \vee a\langle Z \rangle$  and thus  $[\langle Z \rangle] < [\langle \mathbf{1} \rangle]$  in  $\mathbf{BL}_{\mathcal{T}}/L(a\langle Z \rangle)$ .

In the  $p$ -local stable homotopy category, we can take  $\langle Z \rangle = \langle H\mathbb{F}_p \rangle$ . The spectrum  $H\mathbb{F}_p$  is a ring spectrum, and Bousfield [Bou79a] shows that the Bousfield class of any ring spectrum is in the distributive lattice. Let  $IS^0$  be the Brown-Comenetz dual of the sphere. Then Lemma 7.1 in [HP99] shows that  $IS^0 \wedge H\mathbb{F}_p = 0$ , and  $\langle IS^0 \rangle \leq \langle H\mathbb{F}_p \rangle$  so  $\langle IS^0 \rangle \wedge a\langle H\mathbb{F}_p \rangle = \langle 0 \rangle$  by Lemma 2.5 above. This shows that  $IS^0 \in \langle H\mathbb{F}_p \rangle \vee a\langle H\mathbb{F}_p \rangle$ , and hence  $\langle H\mathbb{F}_p \rangle \vee a\langle H\mathbb{F}_p \rangle < \langle S^0 \rangle$ .

In the category  $D(\Lambda)$  of Example 2.16, we can take  $\langle Z \rangle = \langle k \rangle$ . The class  $\langle k \rangle$  is in  $\mathbf{DL}_{\Lambda}$  because  $k$  is a ring object. The dual  $I\Lambda$  of  $\Lambda$  has  $I\Lambda \wedge k = 0$  [DP08, Cor. 4.12] and  $\langle I\Lambda \rangle \leq \langle k \rangle$  [DP08, Lemma 4.8], so  $\langle I\Lambda \rangle \wedge a\langle k \rangle = \langle 0 \rangle$ . Thus we have that  $I\Lambda \in \langle k \rangle \vee a\langle k \rangle < \langle \Lambda \rangle$ .  $\square$

#### 4. RING MAPS AND THE BOUSFIELD LATTICE

**4.1. Ring maps and derived categories.** In this section, we'll establish basic facts about morphisms of Bousfield lattices induced by ring maps, laying the groundwork for the results in Section 5.

**WARNING:** The results of this section hold in an ungraded or a graded setting, and we will be ambiguous with notation. Thus let  $f : R \rightarrow S$  be either a ring homomorphism between two commutative rings, or a graded ring homomorphism between two graded-commutative rings. Let  $\mathbf{Mod}\text{-}R$  denote either the category of right  $R$ -modules, or the category of right graded  $R$ -modules, and let  $D(R)$  denote either the unbounded derived category of  $R$ -modules, or the unbounded derived category of graded  $R$ -modules; in the latter case, the objects of  $D(R)$  are bigraded.

In either case we will use the standard model structure on the category  $Ch(R) = Ch(\mathbf{Mod}\text{-}R)$  of unbounded chain complexes. The weak equivalences are quasi-isomorphisms, the fibrations are dimensionwise surjections, and the cofibrations are dimensionwise injections with cofibrant cokernels. The cofibrant objects are the complexes that can be written as an increasing union of subcomplexes such that the associated quotients are complexes of projectives with zero differentials. Every object is fibrant. See [Hov99, §2.3] or [HPS97, §9.3] for more details.

A ring map  $f : R \rightarrow S$  induces a functor on module categories  $f_* : \mathbf{Mod}\text{-}R \rightarrow \mathbf{Mod}\text{-}S$ , via extension of scalars, where  $f_*(M) = M \otimes_R S$ . This induces a functor  $f_* : Ch(R) \rightarrow Ch(S)$  on chain complexes. The forgetful functor  $f^* : \mathbf{Mod}\text{-}S \rightarrow \mathbf{Mod}\text{-}R$  induces a functor  $f^* : Ch(S) \rightarrow Ch(R)$ , and  $f_*$  and  $f^*$  are adjoints.

**Definition 4.1.** Let  $f_\bullet$  be the left derived functor  $f_\bullet = Lf_* = L(- \otimes_R S) : D(R) \rightarrow D(S)$ . Let  $f^\bullet = Rf^* : D(S) \rightarrow D(R)$  be the right derived functor of the forgetful functor.

**Lemma 4.2.** *The derived functors  $f_\bullet$  and  $f^\bullet$  exist and form an adjoint pair;  $f_\bullet$  is the left adjoint and  $f^\bullet$  is the right adjoint.*

**Proof.** Since  $f_*$  is left adjoint to  $f^*$ , by [DS95, Rmk. 9.8] it suffices to show that  $f^*$  preserves fibrations and trivial fibrations. Since fibrations are degreewise surjections, this is immediate.  $\square$

The functor  $f_\bullet$  is exact (i.e. sends exact triangles to exact triangles), has  $f_\bullet(R) = S$ , and  $f_\bullet(X \wedge Y) = f_\bullet X \wedge f_\bullet Y$  (see [HPS97, Thm. 9.3.1] and note that they consider both the ungraded and graded settings). Since it is a left adjoint, it commutes with coproducts. Since every object is fibrant, we have  $f^\bullet(X) = f^*(X)$  for all  $X$ , so  $f^\bullet$  is exact and commutes with coproducts and products.

**Remark 4.3.** Take  $z \in R = [R, R]_*$ , and consider the morphism  $R \xrightarrow{z} R$  in  $D(R)$ . Applying  $f_\bullet$  to this, we get

$$\begin{aligned} \left( f_\bullet(R) \xrightarrow{f_\bullet(z)} f_\bullet(R) \right) &= \left( R \otimes_R S \xrightarrow{z \otimes 1} R \otimes_R S \right) \\ &= \left( R \otimes_R S \xrightarrow{1 \otimes f(z)} R \otimes_R S \right) = \left( S \xrightarrow{f(z)} S \right). \end{aligned}$$

The following lemma, called the projection formula and proved in [Wei94] for bounded-below complexes, will be used frequently.

**Lemma 4.4.** *(Projection Formula) For all objects  $A$  in  $D(R)$  and  $B$  in  $D(S)$ , we have*

$$f^\bullet(f_\bullet A \wedge B) = A \wedge f^\bullet B.$$

**Proof.** Recall that we can compute the derived tensor product  $- \wedge -$  by taking a cofibrant replacement in either factor. Let  $QX$  represent a choice of cofibrant replacement for a complex  $X$ .

Since every object is fibrant, we have

$$f^\bullet(f_\bullet A \wedge B) = f^*(f_\bullet A \wedge B) = f^*(Q(f_\bullet A) \otimes_S B).$$

To compute  $f_\bullet A$  we use a cofibrant replacement  $QA$ . Since  $f_\bullet R = S$ , and  $f_\bullet$  is exact, and  $f_\bullet$  commutes with coproducts, we see that  $f_\bullet$  takes cofibrant objects to cofibrant objects. Thus  $Q(f_\bullet A) = Q(f_\bullet(QA)) = f_\bullet(QA) = f_*(QA)$ .

At the module level, for  $M \in \mathbf{Mod}\text{-}R$  and  $N \in \mathbf{Mod}\text{-}S$ , we have

$$f^*(f_*(M) \otimes_S N) = f^*((M \otimes_R S) \otimes_S N) = M \otimes_R f^*(N),$$

and this extends to the level of chain complexes, to give

$$f^\bullet(f_\bullet A \wedge B) = f^*(f_*(QA) \otimes_S B) = QA \otimes_R f^*(B) = A \wedge f^\bullet B.$$

□

**Corollary 4.5.** *For all objects  $A$  in  $D(R)$  and  $B$  in  $D(S)$ ,*

$$f_\bullet A \wedge B = 0 \text{ if and only if } A \wedge f^\bullet B = 0.$$

**4.2. Induced maps on Bousfield lattices.** Here we show that the functors  $f_\bullet$  and  $f^\bullet$  induce maps between the Bousfield lattices of  $D(R)$  and  $D(S)$ . If we consider a Bousfield class  $\langle X \rangle$  as the localizing subcategory of  $X$ -acyclics, then we can map this to  $f_\bullet(\langle X \rangle)$  as a subcollection in  $D(S)$ ; however, in general  $f_\bullet(\langle X \rangle)$  will not be triangulated. Instead we make the following definitions.

**Definition 4.6.** Define a map  $f_\bullet : \mathbf{BL}_R \rightarrow \mathbf{BL}_S$  by  $\langle X \rangle \mapsto \langle f_\bullet X \rangle$ . Also, define a map  $f^\bullet : \mathbf{BL}_S \rightarrow \mathbf{BL}_R$  by  $\langle X \rangle \mapsto \langle f^\bullet X \rangle$ . For the rest of this document,  $f_\bullet \langle X \rangle$  and  $f^\bullet \langle X \rangle$  will mean  $\langle f_\bullet X \rangle$  and  $\langle f^\bullet X \rangle$ .

**Proposition 4.7.** *Both  $f_\bullet$  and  $f^\bullet$  induce morphisms on Bousfield lattices that commute with arbitrary joins.*

**Proof.** First we show that  $\langle Y \rangle \leq \langle X \rangle$  implies  $\langle f^\bullet Y \rangle \leq \langle f^\bullet X \rangle$ . Suppose  $\langle Y \rangle \leq \langle X \rangle$  and  $W \wedge f^\bullet X = 0$ . Then Corollary 4.5 implies  $f_\bullet W \wedge X = 0$ . Thus  $f_\bullet W \wedge Y = 0$ , and  $W \wedge f^\bullet Y = 0$ .

This implies that if  $\langle Y \rangle = \langle X \rangle$ , then  $\langle f^\bullet Y \rangle = \langle f^\bullet X \rangle$ , so  $f^\bullet$  is well-defined and order-preserving.

Now suppose  $\langle Y \rangle \leq \langle X \rangle$  and  $f_\bullet X \wedge W = 0$ . Then from Corollary 4.5,  $X \wedge f^\bullet W = 0$ , so  $Y \wedge f^\bullet W = 0$ , which implies  $f_\bullet Y \wedge W = 0$ . Therefore  $f_\bullet$  is order-preserving and well-defined. Both  $f_\bullet$  and  $f^\bullet$  commute with coproducts on the object level, hence with arbitrary joins at the level of Bousfield classes. □

Note that  $f_\bullet$  commutes with the tensor operation,  $f_\bullet(\langle X \wedge Y \rangle) = \langle f_\bullet X \rangle \wedge \langle f_\bullet Y \rangle$ , but in general  $f^\bullet$  does not. See Lemma 4.13 however.

Recall from Section 2 that since  $R \in D(R)$ ,  $\langle \text{Max} \rangle = \langle R \rangle$  in  $\mathbf{BL}_R$ , and every complemented class in  $\mathbf{BL}_R$  is in  $\mathbf{DL}_R$ . Furthermore, complements are unique, and are given by the complementation operator  $a(-)$ .

**Lemma 4.8.** *The functor  $f_\bullet$  maps  $\mathbf{DL}_R$  into  $\mathbf{DL}_S$ , and  $\mathbf{BA}_R$  into  $\mathbf{BA}_S$ . If  $\langle X \rangle$  in  $\mathbf{BA}_R$  has complement  $\langle X^c \rangle$ , then  $\langle f_\bullet X \rangle$  has complement  $\langle f_\bullet(X^c) \rangle$ .*

**Proof.** If  $\langle Y \rangle = \langle Y \wedge Y \rangle$ , then  $\langle f_\bullet Y \rangle = \langle f_\bullet Y \wedge f_\bullet Y \rangle$ .

If  $\langle X \rangle$  has  $\langle X \rangle \vee \langle X^c \rangle = \langle R \rangle$  and  $\langle X \rangle \wedge \langle X^c \rangle = \langle 0 \rangle$ , then  $\langle f_\bullet X \rangle \vee \langle f_\bullet (X^c) \rangle = \langle f_\bullet R \rangle = \langle S \rangle$  and  $\langle f_\bullet X \rangle \wedge \langle f_\bullet (X^c) \rangle = \langle 0 \rangle$ , so  $\langle (f_\bullet X)^c \rangle = \langle f_\bullet (X^c) \rangle$ .  $\square$

We will strengthen and extend this lemma in the next subsection, under additional hypotheses. Next we describe a useful quotient of  $\mathbf{BL}_R$ .

**Definition 4.9.** Fix  $J_f$  to be the image of  $\text{Ker } f_\bullet$  in  $\mathbf{BL}_R$ , in other words  $J_f = \{ \langle X \rangle \mid f_\bullet \langle X \rangle = \langle 0 \rangle \}$ . Also define

$$\langle M_f \rangle := \bigvee_{\langle Y \rangle \in J_f} \langle Y \rangle.$$

**Proposition 4.10.** *The subposet  $J_f$  is a principal ideal in  $\mathbf{BL}_R$  with  $J_f = \langle M_f \rangle \downarrow$ , and  $f_\bullet$  induces a lattice morphism that preserves arbitrary joins,*

$$\overline{f_\bullet} : \mathbf{BL}_R / J_f \rightarrow \mathbf{BL}_S,$$

where  $\overline{f_\bullet} \langle X \rangle = \langle f_\bullet X \rangle$ .

**Proof.** Suppose  $\langle Y \rangle \leq \langle X \rangle$  and  $\langle f_\bullet X \rangle = \langle 0 \rangle$ . Then  $\langle f_\bullet Y \rangle \leq \langle f_\bullet X \rangle$ , so  $\langle f_\bullet Y \rangle = \langle 0 \rangle$  and  $J_f$  is a lattice ideal. Every  $\langle X \rangle$  in  $J_f$  has  $\langle X \rangle \leq \langle M_f \rangle$ , so  $J_f \subseteq \langle M_f \rangle \downarrow$ . And since  $f_\bullet \langle M_f \rangle = 0$ , if  $\langle X \rangle \leq \langle M_f \rangle$  then  $\langle X \rangle \in J_f$ . Therefore  $J_f = \langle M_f \rangle \downarrow$  is principal.

To get an induced map on the quotient lattice, we need to know that if  $[\langle X \rangle] = [\langle Y \rangle]$ , then  $f_\bullet \langle X \rangle = f_\bullet \langle Y \rangle$ . Since  $\langle X \rangle$  and  $\langle Y \rangle$  are equivalent if and only if  $\langle X \rangle \vee \langle M_f \rangle = \langle Y \rangle \vee \langle M_f \rangle$ , and  $\langle f_\bullet M_f \rangle = \langle 0 \rangle$ , we get

$$\langle f_\bullet X \rangle = \langle f_\bullet X \rangle \vee \langle f_\bullet M_f \rangle = f_\bullet (\langle X \rangle \vee \langle M_f \rangle) = \langle f_\bullet Y \rangle \vee \langle f_\bullet M_f \rangle = \langle f_\bullet Y \rangle.$$

Thus  $\overline{f_\bullet}$  is well-defined. It is order and join-preserving since  $f_\bullet$  is.  $\square$

**Remark 4.11.** Note that for any object  $X$  in  $D(R)$ , the projection formula implies  $f_\bullet X = 0$  if and only if  $X \wedge f^\bullet S = 0$ , which is true if and only if  $\langle X \rangle \leq \langle M_f \rangle$ . Therefore by definition  $\langle M_f \rangle = a \langle f^\bullet S \rangle$ , and thus also  $\langle f^\bullet S \rangle = a \langle M_f \rangle$ .

**Proposition 4.12.** *If  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$ , then the quotient functor  $\pi : D(R) \rightarrow D(R) / \langle f^\bullet S \rangle$  induces a lattice isomorphism*

$$\overline{\pi} : \mathbf{BL}_R / J_f \longrightarrow \mathbf{BL}(D(R) / \langle f^\bullet S \rangle).$$

**Proof.** This follows easily from Remark 4.11 and Corollary 3.3, and the fact that  $J_f = \langle M_f \rangle \downarrow$ .  $\square$

We don't have a general criteria for when to expect  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$  to hold. It holds when  $\mathbf{BL}_R = \mathbf{DL}_R$ , thanks to Corollary 2.7. This is the case if  $R$  is Noetherian, for example. On the other hand, consider the ring  $\Lambda$  from Example 2.16, and let  $f : \Lambda \rightarrow k$  be projection onto the degree zero piece. Then  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle k \rangle \vee a \langle k \rangle$ , and as shown in the proof of Proposition 3.5, this is strictly less than  $\langle \Lambda \rangle$ .

**4.3. Maps  $f : R \rightarrow S$  satisfying  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $X$ .** In this subsection we assume the map  $f : R \rightarrow S$  satisfies  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $X$ . In Section 5 we show that this condition holds for the specific map of non-Noetherian rings  $g_\bullet : \Lambda_{\mathbb{Z}(p)} \rightarrow \Lambda_{\mathbb{F}_p}$ , but fails for  $h_\bullet : \Lambda_{\mathbb{Z}(p)} \rightarrow \Lambda_{\mathbb{Q}}$ .

**Lemma 4.13.** *The following are equivalent:*

- (1)  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ ,
- (2)  $f^\bullet W \wedge f^\bullet X = 0$  if and only if  $f^\bullet (W \wedge X) = 0$ ,

$$(3) \ f^\bullet \langle Y \wedge X \rangle = \langle f^\bullet Y \rangle \wedge \langle f^\bullet X \rangle.$$

**Proof.** For (1)  $\Leftrightarrow$  (2), note that  $W \wedge f_\bullet f^\bullet X = 0$  iff  $f^\bullet W \wedge f^\bullet X = 0$ , and  $W \wedge X = 0$  iff  $f^\bullet(W \wedge X) = 0$ .

For (1)  $\Leftrightarrow$  (3), note that  $W \wedge f^\bullet(Y \wedge X) = 0$  iff  $f_\bullet W \wedge (Y \wedge X) = 0$  iff  $(f_\bullet W \wedge Y) \wedge X = 0$ , and  $W \wedge f^\bullet X \wedge f^\bullet Y = 0$  iff  $(f_\bullet W \wedge f_\bullet f^\bullet X) \wedge Y = 0$  iff  $(f_\bullet W \wedge Y) \wedge (f_\bullet f^\bullet X) = 0$ .  $\square$

This is a good setting in which to consider the behavior of the sub-posets  $\mathbf{BA}$  and  $\mathbf{DL}$  under  $f_\bullet$  and  $f^\bullet$ .

**Lemma 4.14.** *Suppose  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ . Then the map  $f^\bullet$  sends  $\mathbf{BA}_S$  into  $\mathbf{BA}_R$  if and only if  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$ . If this is the case, and  $\langle X \rangle \in \mathbf{BA}_S$  has complement  $\langle X^c \rangle$ , then  $\langle f^\bullet X \rangle \in \mathbf{BA}_R$  has complement  $\langle f^\bullet(X^c) \rangle \vee \langle M_f \rangle$ .*

**Proof.** If  $f^\bullet$  injects  $\mathbf{BA}_S$  into  $\mathbf{BA}_R$ , then since  $\langle 0 \rangle$  and  $\langle S \rangle$  are a complemented pair in  $\mathbf{BA}_S$ , the class  $\langle f^\bullet S \rangle$  is complemented in  $\mathbf{BA}_R$ . Its complement must be  $a\langle f^\bullet S \rangle$ , which is  $\langle M_f \rangle$  by Remark 4.11.

For the converse, suppose that  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$ . Now suppose  $\langle X \rangle \in \mathbf{BA}_{D(S)}$ , so  $\langle X \rangle \vee \langle X^c \rangle = \langle S \rangle$  and  $\langle X \rangle \wedge \langle X^c \rangle = \langle 0 \rangle$ . This implies  $\langle f^\bullet X \rangle \vee \langle f^\bullet(X^c) \rangle = \langle f^\bullet S \rangle$  and  $\langle f^\bullet X \rangle \wedge \langle f^\bullet(X^c) \rangle = \langle 0 \rangle$ , using Lemma 4.13.

We calculate that

$$\langle f^\bullet X \rangle \vee (\langle f^\bullet(X^c) \rangle \vee \langle M_f \rangle) = \langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle.$$

Also, we have

$$\begin{aligned} \langle f^\bullet X \rangle \wedge (\langle f^\bullet(X^c) \rangle \vee \langle M_f \rangle) &= (\langle f^\bullet X \rangle \wedge \langle f^\bullet(X^c) \rangle) \vee (\langle f^\bullet X \rangle \wedge \langle M_f \rangle) \\ &= \langle 0 \rangle \vee (\langle f^\bullet X \rangle \wedge \langle M_f \rangle) = \langle 0 \rangle. \end{aligned}$$

The last equality follows from the fact that  $f^\bullet$  is order preserving and  $\langle X \rangle \leq \langle S \rangle$  for all  $\langle X \rangle$ , so  $\langle f^\bullet S \rangle \wedge \langle M_f \rangle = \langle 0 \rangle$  implies  $\langle f^\bullet X \rangle \wedge \langle M_f \rangle = \langle 0 \rangle$  for all  $X$  in  $D(S)$ .

This shows that the complement of  $\langle f^\bullet X \rangle$  is  $\langle f^\bullet(X^c) \rangle \vee \langle M_f \rangle$ .  $\square$

**Proposition 4.15.** *Suppose  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ . The following hold.*

- (1) *The map  $f_\bullet$  sends  $\mathbf{DL}_R$  onto  $\mathbf{DL}_S$ , and the map  $f^\bullet$  injects  $\mathbf{DL}_S$  into  $\mathbf{DL}_R$ .*
- (2) *The map  $f_\bullet$  sends  $\mathbf{BA}_R$  onto  $\mathbf{BA}_S$ , and if  $\langle f^\bullet S \rangle \vee \langle M_f \rangle = \langle R \rangle$  then  $f^\bullet$  injects  $\mathbf{BA}_S$  into  $\mathbf{BA}_R$ .*

**Proof.** Lemma 4.13 implies that if  $\langle Y \rangle = \langle Y \wedge Y \rangle$ , then  $\langle f^\bullet Y \rangle = \langle f^\bullet Y \wedge f^\bullet Y \rangle$ , so  $f^\bullet$  sends  $\mathbf{DL}_S$  to  $\mathbf{DL}_R$  and is injective by hypothesis. The rest follows from Proposition 4.8, Lemma 4.14, and the fact that  $f_\bullet$  is surjective and  $f^\bullet$  is injective.  $\square$

This is also a good setting in which to consider poset adjoints. As a poset map, because  $f^\bullet$  preserves joins on  $\mathbf{BL}_S$ , it has a poset map right adjoint  $r : \mathbf{BL}_R \rightarrow \mathbf{BL}_S$ , see [HP99, Lemma 3.5]. We know

$$r\langle Y \rangle = \bigvee \{ \langle X \rangle \mid f^\bullet \langle X \rangle \leq \langle Y \rangle \}, \text{ and}$$

$$f^\bullet \langle X \rangle \leq \langle Y \rangle \text{ if and only if } \langle X \rangle \leq r\langle Y \rangle.$$

**Proposition 4.16.** *If  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ , then  $f_\bullet \langle X \rangle = r\langle X \rangle$  for all  $\langle X \rangle$ , so*

$$\langle f^\bullet X \rangle \leq \langle Y \rangle \text{ if and only if } \langle X \rangle \leq \langle f_\bullet Y \rangle.$$

**Proof.** First suppose that  $\langle f^\bullet X \rangle \leq \langle Y \rangle$  and  $W \wedge f_\bullet Y = 0$  for some  $W$ . Then Corollary 4.5 implies  $f^\bullet W \wedge Y = 0$ , so  $f^\bullet W \wedge f^\bullet X = 0$ . It follows from Lemma 4.13 that  $f^\bullet(W \wedge X) = 0$ , so  $W \wedge X = 0$ .

For the other direction, if  $\langle X \rangle \leq \langle f_\bullet Y \rangle$ , then  $\langle f^\bullet X \rangle \leq \langle f^\bullet f_\bullet Y \rangle$ . In general we always have  $\langle f^\bullet f_\bullet Y \rangle \leq \langle Y \rangle$ . Indeed, an object  $W$  has  $W \wedge f^\bullet f_\bullet Y = 0$  iff  $f_\bullet W \wedge f_\bullet Y = 0$  iff  $f_\bullet(W \wedge Y) = 0$ , so  $W \wedge Y = 0$  implies  $W \wedge (f^\bullet f_\bullet Y) = 0$ .

This immediately implies that  $\bigvee \{ \langle X \rangle \mid f^\bullet \langle X \rangle \leq \langle Y \rangle \} \leq f_\bullet \langle Y \rangle$ . And the fact that  $f^\bullet \langle f_\bullet Y \rangle \leq \langle Y \rangle$  gives  $\langle f_\bullet Y \rangle \leq \bigvee \{ \langle X \rangle \mid f^\bullet \langle X \rangle \leq \langle Y \rangle \}$ .  $\square$

The BL operation  $f_\bullet$  also preserves arbitrary joins, so has a poset map right adjoint. On the object level, we know that  $f^\bullet$  is right adjoint to  $f_\bullet$ , and so it is natural to ask if  $f^\bullet$  is the right poset adjoint of  $f_\bullet$ .

**Proposition 4.17.** *Assume  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $X$ . Then on the level of Bousfield classes, we have*

$$\langle f_\bullet X \rangle \leq \langle Y \rangle \Leftarrow \langle X \rangle \leq \langle f^\bullet Y \rangle,$$

*but the forward direction need not hold.*

**Proof.** First suppose  $\langle X \rangle \leq \langle f^\bullet Y \rangle$  and  $W \wedge Y = 0$ . Then  $f^\bullet(W \wedge Y) = 0$ , which using Lemma 4.13 means  $f^\bullet W \wedge f^\bullet Y = 0$ , so  $f^\bullet W \wedge X = 0$ , and  $W \wedge f_\bullet X = 0$ .

On the other hand, suppose  $\langle f_\bullet X \rangle \leq \langle Y \rangle$  and  $W \wedge f^\bullet Y = 0$ . Then  $f_\bullet W \wedge Y = 0$ ,  $f_\bullet W \wedge f_\bullet X = 0$ , and  $f_\bullet(W \wedge X) = 0$ . At the BL level, this does not necessarily mean  $W \wedge X = 0$ . (Take, for example,  $Y = 0$ ,  $W = R$ , and  $X$  any object such that  $f_\bullet X = 0$ .)  $\square$

We end this section with another lattice isomorphism.

**Theorem 4.18.** *Suppose  $f_\bullet f^\bullet \langle X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ . There is a lattice isomorphism*

$$\phi : \mathbf{BL}(D(R)/\langle f^\bullet S \rangle) \rightarrow \mathbf{BL}_S,$$

*given by  $\phi \langle X \rangle = \langle f_\bullet \tilde{X} \rangle$ , where  $\pi \tilde{X} = X$ .*

**Proof.** Recall that  $\pi : D(R) \rightarrow D(R)/\langle f^\bullet S \rangle$  is the canonical projection. First we suppose that  $\langle X \rangle, \langle Y \rangle$  in  $\mathbf{BL}(D(R)/\langle f^\bullet S \rangle)$  have  $\langle X \rangle \leq \langle Y \rangle$ , and will show that then  $\phi \langle X \rangle \leq \phi \langle Y \rangle$ . Fix a choice of  $\tilde{X}$  and  $\tilde{Y}$  such that  $\pi \tilde{X} = X$  and  $\pi \tilde{Y} = Y$ .

Take  $W \in D(S)$  such that  $W \wedge f_\bullet \tilde{Y} = 0$ . We wish to show that  $W \wedge f_\bullet \tilde{X} = 0$ . Corollary 4.5 implies that  $f^\bullet W \wedge \tilde{Y} = 0$ . So

$$0 = \pi(f^\bullet W \wedge \tilde{Y}) = \pi f^\bullet W \wedge \pi \tilde{Y} = \pi f^\bullet W \wedge Y.$$

By hypothesis, this implies that  $\pi f^\bullet W \wedge X = 0$ . Thus  $\pi(f^\bullet W \wedge \tilde{X}) = 0$ , and  $f^\bullet W \wedge \tilde{X} \in \langle f^\bullet S \rangle$ , so  $f^\bullet W \wedge \tilde{X} \wedge f^\bullet S = 0$ . Again the projection formula implies that  $f_\bullet(f^\bullet W \wedge \tilde{X}) = 0$ , so  $f_\bullet f^\bullet W \wedge f_\bullet \tilde{X} = 0$ . Since we're assuming  $\langle f_\bullet f^\bullet W \rangle = \langle W \rangle$ , we conclude that  $W \wedge f_\bullet \tilde{X} = 0$ , as desired.

This shows that  $\phi$  is order-preserving. By symmetry, it also shows that  $\phi$  is well-defined, independent of choice of representative or preimage.

The map  $\phi$  is surjective by assumption: given  $\langle Y \rangle \in \mathbf{BL}_S$ , we get  $\phi \langle \pi f^\bullet Y \rangle = \langle f_\bullet f^\bullet Y \rangle = \langle Y \rangle$ .

For injectivity, suppose  $\langle f_\bullet \tilde{X} \rangle = \langle f_\bullet \tilde{Y} \rangle$ . We will show  $\langle \pi \tilde{X} \rangle \leq \langle \pi \tilde{Y} \rangle$ , and injectivity follows by symmetry. Suppose  $W \in D(R)/\langle f^\bullet S \rangle$  has  $W \wedge \pi \tilde{Y} = 0$ . Choose  $\tilde{W}$  so  $\pi \tilde{W} = W$ . Then  $\pi(\tilde{W} \wedge \tilde{Y}) = 0$ , so  $\tilde{W} \wedge \tilde{Y} \wedge f^\bullet S = 0$ , and  $0 = f_\bullet(\tilde{W} \wedge \tilde{Y}) = f_\bullet \tilde{W} \wedge f_\bullet \tilde{Y}$ .



By hypothesis, this implies that  $0 = f_\bullet \tilde{W} \wedge f_\bullet \tilde{X} = f_\bullet (\tilde{W} \wedge \tilde{X})$ , so  $\tilde{W} \wedge \tilde{X} \wedge f^\bullet S = 0$ . This says that  $0 = \pi \tilde{W} \wedge \pi \tilde{X} = W \wedge \pi \tilde{X}$ , and we conclude that  $\phi$  is injective.

The inverse of  $\phi$  is clearly given by  $\phi^{-1}\langle Y \rangle = \langle \pi f^\bullet Y \rangle$ .  $\square$

## 5. NON-NOETHERIAN RINGS

Here we will investigate maps between several graded non-Noetherian rings. **All rings and modules in this section are graded, and objects in derived categories are bigraded.**

**Definition 5.1.**

- (1) Fix a prime  $p$  and integers  $n_i > 1$  for  $i \geq 1$ , and set

$$\Lambda_{\mathbb{Z}_{(p)}} := \frac{\mathbb{Z}_{(p)}[x_1, x_2, \dots]}{(x_1^{n_1}, x_2^{n_2}, \dots)}, \quad \Lambda_{\mathbb{F}_p} := \frac{\mathbb{F}_p[x_1, x_2, \dots]}{(x_1^{n_1}, x_2^{n_2}, \dots)} \quad \text{and} \quad \Lambda_{\mathbb{Q}} := \frac{\mathbb{Q}[x_1, x_2, \dots]}{(x_1^{n_1}, x_2^{n_2}, \dots)}.$$

Grade the  $x_i$  so that  $\Lambda_{\mathbb{Z}_{(p)}}$ ,  $\Lambda_{\mathbb{F}_p}$ , and  $\Lambda_{\mathbb{Q}}$  are graded-connected and finitely-generated in each module degree, for example by setting  $\deg(x_i) = 2^i$ .

- (2) Fix  $g : \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{Z}_{(p)}}/p\Lambda_{\mathbb{Z}_{(p)}} = \Lambda_{\mathbb{F}_p}$  to be the projection map, and fix  $h : \Lambda_{\mathbb{Z}_{(p)}} \hookrightarrow \Lambda_{\mathbb{Q}}$  to be inclusion.
- (3) Let  $g_\bullet : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightarrow D(\Lambda_{\mathbb{F}_p})$  and  $h_\bullet : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightarrow D(\Lambda_{\mathbb{Q}})$  be the induced functors on unbounded derived categories of chain complexes of graded modules, as in Section 2. Let  $g^\bullet$  and  $h^\bullet$  denote their corresponding right adjoints.

**Remark 5.2.** Note that  $g^\bullet \Lambda_{\mathbb{F}_p}$  can be represented in  $D(\Lambda_{\mathbb{Z}_{(p)}})$  by the chain complex

$$\left( \cdots \rightarrow 0 \rightarrow \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{p} \Lambda_{\mathbb{Z}_{(p)}} \rightarrow 0 \rightarrow \cdots \right).$$

Furthermore,  $g^\bullet \Lambda_{\mathbb{F}_p}$  fits into the exact triangle  $\Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{p} \Lambda_{\mathbb{Z}_{(p)}} \rightarrow g^\bullet \Lambda_{\mathbb{F}_p}$  in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ .

**Proposition 5.3.** *For all  $X$  in  $D(\Lambda_{\mathbb{F}_p})$ , we have  $g_\bullet g^\bullet X \cong X \oplus \Sigma X$ . Therefore  $\text{th}(g_\bullet g^\bullet X) = \text{th}(X)$  and  $\langle g_\bullet g^\bullet X \rangle = \langle X \rangle$ .*

**Proof.** Using  $g_\bullet \Lambda_{\mathbb{Z}_{(p)}} = \Lambda_{\mathbb{F}_p}$ , and Remarks 4.3 and 5.2, we see that  $g_\bullet g^\bullet \Lambda_{\mathbb{F}_p}$  in  $D(\Lambda_{\mathbb{F}_p})$  is

$$\left( \cdots \rightarrow 0 \rightarrow \Lambda_{\mathbb{F}_p} \xrightarrow{0} \Lambda_{\mathbb{F}_p} \rightarrow 0 \rightarrow \cdots \right),$$

which is just  $\Lambda_{\mathbb{F}_p} \oplus \Sigma \Lambda_{\mathbb{F}_p}$ .

Now let  $X$  be a cofibrant representative for an arbitrary element of  $D(\Lambda_{\mathbb{F}_p})$ . Cofibrant complexes are object-wise projective. Since  $\Lambda_{\mathbb{F}_p}$  is a local ring, projectives are free. Thus  $X$  has the form

$$\cdots \rightarrow \coprod_{I_3} \Lambda_{\mathbb{F}_p} \xrightarrow{d_2} \coprod_{I_2} \Lambda_{\mathbb{F}_p} \xrightarrow{d_1} \coprod_{I_1} \Lambda_{\mathbb{F}_p} \xrightarrow{d_0} \coprod_{I_0} \Lambda_{\mathbb{F}_p} \rightarrow \cdots.$$

Each differential is a direct sum of maps  $\Lambda_{\mathbb{F}_p} \rightarrow \Lambda_{\mathbb{F}_p}$ , which we can think of as elements of  $\Lambda_{\mathbb{F}_p}$ .

Since every object is fibrant,  $g^\bullet X = g^* X$ , and this is the complex

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \coprod_{I_2} \Lambda_{\mathbb{Z}_{(p)}} & \xrightarrow{\overline{d_1}} & \coprod_{I_1} \Lambda_{\mathbb{Z}_{(p)}} & \xrightarrow{\overline{d_0}} & \coprod_{I_0} \Lambda_{\mathbb{Z}_{(p)}} \longrightarrow \cdots \\ & & \oplus & \searrow \oplus p & \oplus & \searrow \oplus p & \oplus & \searrow \oplus p \\ \cdots & \longrightarrow & \coprod_{I_3} \Lambda_{\mathbb{Z}_{(p)}} & \xrightarrow{\overline{d_2}} & \coprod_{I_2} \Lambda_{\mathbb{Z}_{(p)}} & \xrightarrow{\overline{d_1}} & \coprod_{I_1} \Lambda_{\mathbb{Z}_{(p)}} & \xrightarrow{\overline{d_0}} & \coprod_{I_0} \Lambda_{\mathbb{Z}_{(p)}} \longrightarrow \cdots \end{array}$$

Here  $\overline{d_i}$  is a direct sum of maps  $\Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{Z}_{(p)}}$  that correspond to preimages via  $g : \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{F}_p}$  of the elements of  $\Lambda_{\mathbb{F}_p}$  comprising  $d_i$ . We claim that this complex is cofibrant. First note that it is the cofiber in  $Ch(\Lambda_{\mathbb{Z}_{(p)}})$  of  $W \xrightarrow{p} W$ , where  $W$  is the complex

$$\cdots \rightarrow \coprod_{I_3} \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{\overline{d_2}} \coprod_{I_2} \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{\overline{d_1}} \coprod_{I_1} \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{\overline{d_0}} \coprod_{I_0} \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \cdots$$

Since  $X$  is cofibrant, so is  $W$ . For  $X$  is an increasing union of complexes, such that the associated quotients are complexes of free  $\Lambda_{\mathbb{F}_p}$ -modules with zero differentials. By replacing each  $\Lambda_{\mathbb{F}_p}$  with  $\Lambda_{\mathbb{Z}_{(p)}}$  and each map, thought of as an element of  $\Lambda_{\mathbb{F}_p}$ , with its preimage via  $g$  (in a compatible way), we construct  $W$  as such an increasing union. Since  $W$  is cofibrant, so is  $g^* X$ .

Therefore we can compute  $g_\bullet(g^\bullet X) = g_\bullet(g^* X) = g_*(g^* X)$ . But  $g_*(W) = X$ , and  $g(p) = 0$ , so this gives  $g_\bullet(g^\bullet X) = X \oplus \Sigma X$ .  $\square$

This proposition allows us to apply all the results of Section 4.3 to the case  $g_\bullet : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightarrow D(\Lambda_{\mathbb{F}_p})$ .

Next, we point out an important difference between  $D(\Lambda_{\mathbb{Z}_{(p)}})$  and  $D(\Lambda_{\mathbb{F}_p})$ . Recall from Example 2.16 that  $\text{BA}(\Lambda_{\mathbb{F}_p})$  is trivial, and the module  $\langle I\Lambda_{\mathbb{F}_p} \rangle$  is a minimum nonzero Bousfield class. The latter fact plays a significant role in [DP08].

Recall that, given a self-map  $X \xrightarrow{f} X$  in any derived category  $D(R)$ , the telescope  $f^{-1}X$  is defined to be the cofiber of the map  $\coprod_{i \geq 0} X_i \xrightarrow{1-f} \coprod_{i \geq 0} X_i$ , where  $X_i = X$  for all  $i$  and the map sends each summand  $X_i \rightarrow X_i \coprod X_{i+1}$  by  $(1-f)(x) = (x, -f(x))$ . This is a minimal weak colimit (see e.g. [HPS97, Prop. 2.2.4]), so for all  $n$  we have

$$H_n(f^{-1}X) \cong \varinjlim (H_n(X) \xrightarrow{H(f)} H_n(X) \rightarrow \cdots).$$

**Proposition 5.4.** *The classes  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$  and  $\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$  form a complemented pair in  $\text{BA}(\Lambda_{\mathbb{Z}_{(p)}})$ . Thus there is no minimum nonzero Bousfield class in  $\text{BL}(\Lambda_{\mathbb{Z}_{(p)}})$ .*

**Proof.** Consider the self-map  $\Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{p} \Lambda_{\mathbb{Z}_{(p)}}$  in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ . The telescope  $p^{-1}\Lambda_{\mathbb{Z}_{(p)}}$  is quasi-isomorphic to a module concentrated in chain degree zero, with zeroth homology the  $\Lambda_{\mathbb{Z}_{(p)}}$ -module

$$\varinjlim \left( \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{p} \Lambda_{\mathbb{Z}_{(p)}} \rightarrow \cdots \right) \cong \frac{\mathbb{Q}[x_1, x_2, \dots]}{(x_1^{n_1}, x_2^{n_2}, \dots)} = \Lambda_{\mathbb{Q}}.$$

Thus we can identify  $p^{-1}\Lambda_{\mathbb{Z}_{(p)}}$  with  $h^\bullet \Lambda_{\mathbb{Q}}$ . As noted above, the cofiber of the map  $\Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{p} \Lambda_{\mathbb{Z}_{(p)}}$  is  $g^\bullet \Lambda_{\mathbb{F}_p}$ . It is then not hard to see that  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle \vee \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle \Lambda_{\mathbb{Z}_{(p)}} \rangle$ .

To compute  $g^\bullet \Lambda_{\mathbb{F}_p} \wedge h^\bullet \Lambda_{\mathbb{Q}}$ , we use the chain complex description of  $g^\bullet \Lambda_{\mathbb{F}_p}$  given in Remark 5.2, and find that  $g^\bullet \Lambda_{\mathbb{F}_p} \wedge h^\bullet \Lambda_{\mathbb{Q}}$  is represented by

$$\left( \cdots \rightarrow 0 \rightarrow \Lambda_{\mathbb{Q}} \xrightarrow{p} \Lambda_{\mathbb{Q}} \rightarrow 0 \rightarrow \cdots \right),$$

which is zero in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ .

Therefore the classes  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$  and  $\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$  form a nontrivial complemented pair in  $\text{BA}(\Lambda_{\mathbb{Z}_{(p)}})$ . Suppose  $\langle Z \rangle$  were a minimum nonzero Bousfield class. Then  $\langle Z \rangle \leq \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ ,  $\langle Z \rangle \leq \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ , and  $g^\bullet \Lambda_{\mathbb{F}_p} \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$  imply  $\langle Z \rangle \wedge \langle g^\bullet \Lambda_{\mathbb{F}_p} \vee h^\bullet \Lambda_{\mathbb{Q}} \rangle = 0$ . This would force  $Z = 0$ .  $\square$

The subcategory  $\text{th}(g^\bullet \Lambda_{\mathbb{F}_p})$  is a thick subcategory of compact objects in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ . It is clearly nonzero, and the inclusion  $\text{th}(g^\bullet \Lambda_{\mathbb{F}_p}) \subseteq \text{th}(\Lambda_{\mathbb{Z}_{(p)}})$  is proper. Indeed, if  $\text{th}(g^\bullet \Lambda_{\mathbb{F}_p}) = \text{th}(\Lambda_{\mathbb{Z}_{(p)}})$  then we would have  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle \Lambda_{\mathbb{Z}_{(p)}} \rangle$ , which contradicts  $g^\bullet \Lambda_{\mathbb{F}_p} \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ .

**Definition 5.5.** Let  $L : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightarrow D(\Lambda_{\mathbb{Z}_{(p)}})$  be finite localization at  $\text{th}(g^\bullet \Lambda_{\mathbb{F}_p})$ . Let  $C$  denote the corresponding colocalization; thus for each  $X$  there is an exact triangle  $CX \rightarrow X \rightarrow LX$ .

See [HPS97, Ch. 3] or [Kra10] for a discussion of Bousfield localization. Recall that we say an object  $X$  is  $L$ -acyclic if  $L(X) = 0$ , and  $L$ -local if it is in the essential image of  $L$ .

**Definition 5.6.** The inclusion  $\mathbb{Z}_{(p)} \hookrightarrow \mathbb{Q}$  induces a morphism  $\Lambda_{\mathbb{Z}_{(p)}} \rightarrow h^\bullet \Lambda_{\mathbb{Q}}$  in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ . Let  $F$  be the fiber of this map, so  $F \rightarrow \Lambda_{\mathbb{Z}_{(p)}} \rightarrow h^\bullet \Lambda_{\mathbb{Q}}$  is an exact triangle in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ .

**Lemma 5.7.** In  $D(\Lambda_{\mathbb{Z}_{(p)}})$ , the object  $\Sigma F$  is quasi-isomorphic to the  $\Lambda_{\mathbb{Z}_{(p)}}$ -module  $\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}$  concentrated in degree zero.

**Proof.** This is a straightforward calculation using the long exact sequence in homology.  $\square$

**Proposition 5.8.** The localization functor  $L$  is smashing, with  $L\Lambda_{\mathbb{Z}_{(p)}} = h^\bullet \Lambda_{\mathbb{Q}}$  and  $C\Lambda_{\mathbb{Z}_{(p)}} = F$ . It has the following acyclics and locals.

$$\begin{aligned} L\text{-acyclics} &= \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p}) = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle L\Lambda_{\mathbb{Z}_{(p)}} \rangle = \langle M_g \rangle, \\ L\text{-locals} &= \text{loc}(h^\bullet \Lambda_{\mathbb{Q}}) = \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle C\Lambda_{\mathbb{Z}_{(p)}} \rangle = \langle M_h \rangle = \langle F \rangle. \end{aligned}$$

**Proof.** All finite localizations are smashing localizations, which means  $LX = L\Lambda_{\mathbb{Z}_{(p)}} \wedge X$ . Thus the  $L$ -acyclics are precisely  $\langle L\Lambda_{\mathbb{Z}_{(p)}} \rangle$ . Finite localization at  $\text{th}(g^\bullet \Lambda_{\mathbb{F}_p})$  means also that the  $L$ -acyclics are  $\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ .

Next we show that the  $L$ -acyclics are the same as  $\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ . Suppose  $X$  is  $L$ -acyclic. Then  $X \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ . Since  $g^\bullet \Lambda_{\mathbb{F}_p} \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ , this implies that  $X \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ . Conversely, suppose that  $X \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ . Then

$$\langle X \wedge g^\bullet \Lambda_{\mathbb{F}_p} \rangle \vee \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle X \wedge \Lambda_{\mathbb{Z}_{(p)}} \rangle = \langle X \rangle,$$

so  $\langle X \wedge g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle X \rangle$  and  $\langle X \rangle \leq \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ . Since  $g^\bullet \Lambda_{\mathbb{F}_p} \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$  is  $L$ -acyclic, we have  $L\Lambda_{\mathbb{Z}_{(p)}} \wedge g^\bullet \Lambda_{\mathbb{F}_p} = 0$ , so  $L\Lambda_{\mathbb{Z}_{(p)}} \wedge X = LX = 0$ , and  $X$  is  $L$ -acyclic.

With any smashing localization, the classes  $\langle L\mathbb{1} \rangle$  and  $\langle C\mathbb{1} \rangle$  are a complemented pair, where  $\mathbb{1}$  is the tensor unit. Furthermore, the  $L$ -locals are precisely  $\langle C\mathbb{1} \rangle$ . Thus in the present context, since  $L : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightarrow D(\Lambda_{\mathbb{Z}_{(p)}})$  is smashing and  $\langle L\Lambda_{\mathbb{Z}_{(p)}} \rangle =$

$\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$  is complemented by  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ , we see that the  $L$ -locals  $\langle C\Lambda_{\mathbb{Z}_{(p)}} \rangle$  are precisely  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ .

From Remark 4.11 we know that  $a\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle M_g \rangle$ . But Proposition 5.4 shows that  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$  is complemented by  $\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ , so  $a\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ . Likewise,  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle M_h \rangle$ .

Next we will show that  $F \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$  in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ . From Lemma 5.7, this is true if and only if  $(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$  in  $D(\Lambda_{\mathbb{Z}_{(p)}})$ . As in the proof of Proposition 5.4, we can identify  $h^\bullet \Lambda_{\mathbb{Q}}$  with the telescope  $p^{-1}\Lambda_{\mathbb{Z}_{(p)}}$ , so in  $D(\Lambda_{\mathbb{Z}_{(p)}})$  there is an exact triangle

$$\bigoplus \Lambda_{\mathbb{Z}_{(p)}} \xrightarrow{1-p} \bigoplus \Lambda_{\mathbb{Z}_{(p)}} \longrightarrow h^\bullet \Lambda_{\mathbb{Q}}.$$

Applying  $(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \wedge -$  to this, we see that  $(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \wedge h^\bullet \Lambda_{\mathbb{Q}}$  is the telescope  $p^{-1}(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}})$ . This has zero homology away from degree zero, and its degree zero homology also vanishes because  $p \in \Lambda_{\mathbb{Z}_{(p)}}$  so the direct limit has all zero maps. Therefore  $F \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ , and  $F$  is  $L$ -acyclic.

From the triangle  $F \rightarrow \Lambda_{\mathbb{Z}_{(p)}} \rightarrow h^\bullet \Lambda_{\mathbb{Q}}$ , we get a triangle

$$L\Lambda_{\mathbb{Z}_{(p)}} \wedge F \longrightarrow L\Lambda_{\mathbb{Z}_{(p)}} \longrightarrow L\Lambda_{\mathbb{Z}_{(p)}} \wedge h^\bullet \Lambda_{\mathbb{Q}}.$$

Since  $L\Lambda_{\mathbb{Z}_{(p)}} \wedge F = 0$ ,  $L\Lambda_{\mathbb{Z}_{(p)}} \cong L\Lambda_{\mathbb{Z}_{(p)}} \wedge h^\bullet \Lambda_{\mathbb{Q}}$ . Since  $h^\bullet \Lambda_{\mathbb{Q}} \in \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ , it is  $L$ -local and  $L\Lambda_{\mathbb{Z}_{(p)}} \wedge h^\bullet \Lambda_{\mathbb{Q}} \cong h^\bullet \Lambda_{\mathbb{Q}}$ . Therefore  $L\Lambda_{\mathbb{Z}_{(p)}} \cong h^\bullet \Lambda_{\mathbb{Q}}$ . The exact triangle  $C\Lambda_{\mathbb{Z}_{(p)}} \rightarrow \Lambda_{\mathbb{Z}_{(p)}} \rightarrow L\Lambda_{\mathbb{Z}_{(p)}}$  then forces  $C\Lambda_{\mathbb{Z}_{(p)}} \cong F$ .

It only remains to show that the  $L$ -locals are given by  $\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})$ . But with a smashing localization, the  $L$ -locals are always form a localizing subcategory, and in addition when  $T = \text{loc}(\mathbf{1})$  we always have  $L\text{-locals} = \text{loc}(L\mathbf{1})$ .  $\square$

The last two propositions show that  $\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle \vee \langle M_g \rangle = \langle \Lambda_{\mathbb{Z}_{(p)}} \rangle$  in  $\text{BL}(\Lambda_{\mathbb{Z}_{(p)}})$ , so Lemma 4.14 and Propositions 4.15 and 4.12 apply in full to this setting.

Recall that  $J_g := \{\langle X \rangle \in \text{BL}(\Lambda_{\mathbb{Z}_{(p)}}) \mid g_\bullet \langle X \rangle = \langle 0 \rangle\}$ . The next theorem shows that the lattice map in Proposition 4.10 becomes an isomorphism.

**Theorem 5.9.** *The functor  $g_\bullet$  induces a lattice isomorphism*

$$\overline{g_\bullet} : \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})/J_g \rightarrow \text{BL}(\Lambda_{\mathbb{F}_p}),$$

*with inverse  $g^\bullet$ .*

**Proof.** Proposition 4.10 showed that  $J_g = \langle M_g \rangle \downarrow = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle \downarrow$  is a principal ideal, and  $\overline{g_\bullet}$  is a lattice morphism. We know  $g^\bullet : \langle Y \rangle \mapsto [\langle g^\bullet Y \rangle]$  is a lattice morphism, and must only check these are inverses.

Proposition 5.3 shows that  $\overline{g_\bullet} g^\bullet \langle X \rangle = \langle g_\bullet g^\bullet X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ .

As noted earlier, we always have  $\langle g^\bullet g_\bullet Y \rangle \leq \langle Y \rangle$  for all  $\langle Y \rangle$ . To prove  $[g^\bullet \overline{g_\bullet} \langle Y \rangle] = [\langle Y \rangle]$ , it remains to show that  $\langle g^\bullet g_\bullet Y \rangle \vee \langle M_g \rangle \geq \langle Y \rangle \vee \langle M_g \rangle$  for all  $\langle Y \rangle$ .

So take  $W$  with  $W \wedge g^\bullet g_\bullet Y = 0$ , and  $W \in \langle M_g \rangle = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ . From the last proposition we get that  $W$  is  $L$ -acyclic, so  $W \wedge L\Lambda_{\mathbb{Z}_{(p)}} = 0$ . But then  $W \wedge Y \wedge L\Lambda_{\mathbb{Z}_{(p)}} = 0$ , so  $W \wedge Y$  is also  $L$ -acyclic.

Now  $W \wedge g^\bullet g_\bullet Y = 0$  implies  $g_\bullet W \wedge g_\bullet Y = g_\bullet(W \wedge Y) = 0$ , so  $W \wedge Y \wedge g^\bullet \Lambda_{\mathbb{F}_p} = 0$ . Using the last proposition, this says that  $W \wedge Y$  is  $L$ -local.

Any object that is both acyclic and local with respect to a localization functor must be zero, because there are no nonzero morphisms from an acyclic to a local object. So we conclude that  $W \wedge Y = 0$ , and therefore  $W \in \langle Y \rangle \wedge \langle M_g \rangle$ .  $\square$

Our next goal is to show that this is actually a splitting of lattices.

**Definition 5.10.**

- (1) Let  $i_g : \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p}) \rightarrow D(\Lambda_{\mathbb{Z}_{(p)}})$  and  $i_h : \text{loc}(h^\bullet \Lambda_{\mathbb{Q}}) \rightarrow D(\Lambda_{\mathbb{Z}_{(p)}})$  denote the inclusions.
- (2) Given a localizing subcategory  $\mathcal{S}$  in a well generated tensor triangulated category  $\mathcal{T}$ , let

$$\text{BL}(\mathcal{T})|_{\mathcal{S}} = \{\langle X \rangle \in \text{BL}_{\mathcal{T}} \mid X \in \mathcal{S}\} \subseteq \text{BL}(\mathcal{T}).$$

But note that  $\langle X \rangle = \langle Y \rangle$  and  $X \in \mathcal{S}$  does not imply  $Y \in \mathcal{S}$  in general.

**Lemma 5.11.** *The inclusions  $i_g$  and  $i_h$  induce lattice morphisms on Bousfield lattices that preserve arbitrary joins.*

$$i_g : \text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})) \rightarrow \text{BL}_{\Lambda_{\mathbb{Z}_{(p)}}}, \text{ where } i_g \langle X \rangle = \langle i_g X \rangle = \langle X \rangle, \text{ and}$$

$$i_h : \text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})) \rightarrow \text{BL}_{\Lambda_{\mathbb{Z}_{(p)}}}, \text{ where } i_h \langle Y \rangle = \langle i_h Y \rangle = \langle Y \rangle.$$

**Proof.** Take  $X, Y \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$  such that  $\langle X \rangle \leq \langle Y \rangle$  in  $\text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p}))$ . Now suppose  $W \in D(\Lambda_{\mathbb{Z}_{(p)}})$  has  $W \wedge Y = 0$ . Then  $W \wedge g^\bullet \Lambda_{\mathbb{F}_p} \wedge Y = 0$ . But  $W \wedge g^\bullet \Lambda_{\mathbb{F}_p} \in \text{loc}(\Lambda_{\mathbb{Z}_{(p)}} \wedge g^\bullet \Lambda_{\mathbb{F}_p}) = \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ , so by hypothesis we have  $W \wedge g^\bullet \Lambda_{\mathbb{F}_p} \wedge X = 0$ . From Proposition 5.8 we conclude that  $W \wedge X$  is  $L$ -local.

Since  $X \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ , it is  $L$ -acyclic, and  $W \wedge X$  is also  $L$ -acyclic. Any object that is both acyclic and local must be zero, so  $W \wedge X = 0$  as desired. Therefore  $i_g$  induces an order-preserving and well-defined map on Bousfield lattices. Coproducts in both  $\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$  and  $D(\Lambda_{\mathbb{Z}_{(p)}})$  are given by degreewise direct sums of modules. So  $i_g$  preserves arbitrary coproducts on the object level, and thus arbitrary joins on the level of Bousfield classes.

A similar argument shows the same for  $i_h$ .  $\square$

This lemma does not generalize to arbitrary localizing subcategory inclusions, but we do have the following lemma, which is easy to prove.

**Lemma 5.12.** *If  $i : \mathcal{S} \rightarrow \mathcal{T}$  is the inclusion of a localizing subcategory, and induces a morphism on Bousfield lattices, then  $\text{BL}(\mathcal{S}) \cong \text{BL}(\mathcal{T})|_{\mathcal{S}}$ .*

**Proposition 5.13.** *The following hold.*

$$\text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})) \cong \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})|_{\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})} = \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle \downarrow, \text{ and}$$

$$\text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})) \cong \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})|_{\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})} = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle \downarrow.$$

**Proof.** The isomorphisms on the left come from Lemmas 5.11 and 5.12. For the equalities on the right, we need to know that for all  $X \in D(\Lambda_{\mathbb{Z}_{(p)}})$  we have  $\langle X \rangle \leq \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$  if and only if  $X \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ , and  $\langle X \rangle \leq \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$  if and only if  $X \in \text{loc}(h^\bullet \Lambda_{\mathbb{Q}})$ . It is always the case that  $X \in \text{loc}(Y)$  implies  $\langle X \rangle \leq \langle Y \rangle$ . If  $\langle X \rangle \leq \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ , then by Proposition 5.8,  $X \in \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \text{loc}(h^\bullet \Lambda_{\mathbb{Q}})$ . Similarly, if  $\langle X \rangle \leq \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ , then  $X \in \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle = \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$ .  $\square$

**Theorem 5.14.** *There is a lattice isomorphism*

$$\Phi : \text{BL}(\Lambda_{\mathbb{Z}_{(p)}}) \xrightarrow{\sim} \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})|_{\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})} \times \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})|_{\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})}, \text{ where}$$

$$\Phi \langle X \rangle = (\langle X \wedge g^\bullet \Lambda_{\mathbb{F}_p} \rangle, \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle).$$

The inverse is given by  $\Phi' : (\langle X \rangle, \langle Y \rangle) \mapsto \langle X \rangle \vee \langle Y \rangle$ .

**Proof.** Note that  $X \wedge Z \in \text{loc}(\Lambda_{\mathbb{Z}_{(p)}} \wedge Z) = \text{loc}(Z)$  for any  $Z$ . It's clear that both  $\Phi$  and  $\Phi'$  are lattice morphisms. We compute  $\Phi'\Phi\langle X \rangle$  as

$$\langle X \wedge g^\bullet \Lambda_{\mathbb{F}_p} \rangle \vee \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle X \rangle \wedge (\langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle \vee \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle) = \langle X \rangle \wedge \langle \Lambda_{\mathbb{Z}_{(p)}} \rangle = \langle X \rangle.$$

On the other hand, Proposition 5.8 shows that, for  $X \in \text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})$  and  $Y \in \text{loc}(h^\bullet \Lambda_{\mathbb{Q}})$ , we can compute  $\Phi\Phi'(\langle X \rangle, \langle Y \rangle)$  as

$$(\langle X \vee Y \rangle \wedge \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle, \langle X \vee Y \rangle \wedge \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle) = (\langle X \rangle, \langle Y \rangle),$$

because  $X$  is  $L$ -acyclic and  $Y$  is  $L$ -local.  $\square$

**Remark 5.15.** As one can see from the proofs, the last four results hold more generally, in the context of any smashing localization functor on any well generated tensor triangulated category  $\mathbb{T}$  with  $\mathbb{T} = \text{loc}(\mathbb{1})$ . If  $K : \mathbb{T} \rightarrow \mathbb{T}$  is a smashing localization functor, with colocalization  $C$ , the statements all hold if we replace  $D(\Lambda_{\mathbb{Z}_{(p)}})$  by  $\mathbb{T}$ , and  $g^\bullet \Lambda_{\mathbb{F}_p}$  by  $C\mathbb{1}$  and  $h^\bullet \Lambda_{\mathbb{Q}}$  by  $K\mathbb{1}$ , since the  $K$ -acyclics  $= \text{loc}(C\mathbb{1}) = \langle K\mathbb{1} \rangle$  and  $K$ -locals  $= \text{loc}(K\mathbb{1}) = \langle C\mathbb{1} \rangle$ . This was essentially shown by Iyengar and Krause in Proposition 6.12 and Lemma 6.13 of [IK11]. The following results are specific to our context.

**Corollary 5.16.** *The functor  $g_\bullet$  induces a lattice isomorphism*

$$g_\bullet : \text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})) \xrightarrow{\sim} \text{BL}(\Lambda_{\mathbb{F}_p}), \text{ with } \langle X \rangle \mapsto \langle g_\bullet X \rangle.$$

**Proof.** This follows by combining Propositions 5.13 and Theorems 5.9 and 5.14, and the fact that  $J_g = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle \downarrow \cong \text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))$ .

$$\text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})) \cong \frac{\text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p})) \times \text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))}{0 \times \text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))} \cong \text{BL}(\Lambda_{\mathbb{Z}_{(p)}})/J_g \xrightarrow{\sim} \text{BL}(\Lambda_{\mathbb{F}_p}).$$

$\square$

**Corollary 5.17.** *The functors  $g_\bullet$  and  $h_\bullet$  induce a lattice isomorphism*

$$\text{BL}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \text{BL}(\Lambda_{\mathbb{F}_p}) \times \text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})), \text{ where}$$

$$\langle X \rangle \mapsto (g_\bullet \langle X \rangle, \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle).$$

*The inverse is given by*

$$(\langle Y \rangle, \langle Z \rangle) \mapsto g^\bullet \langle Y \rangle \vee \langle i_h Z \rangle.$$

**Proof.** This is basically a restatement of Theorem 5.14, along with the observation that

$$g_\bullet \langle X \wedge g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle g_\bullet X \rangle \wedge \langle g_\bullet g^\bullet \Lambda_{\mathbb{F}_p} \rangle = \langle g_\bullet X \rangle \wedge \langle \Lambda_{\mathbb{F}_p} \rangle = \langle g_\bullet X \rangle.$$

$\square$

**Corollary 5.18.** *The isomorphism in Corollary 5.17 induces a splitting of the distributive lattices and Boolean algebras*

$$\text{DL}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \text{DL}(\Lambda_{\mathbb{F}_p}) \times \text{DL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})),$$

$$\text{BA}(\Lambda_{\mathbb{Z}_{(p)}}) \cong \text{BA}(\Lambda_{\mathbb{F}_p}) \times \text{BA}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}})).$$

**Proof.** Much of this follows from Proposition 4.15. First consider the distributive lattice. If  $\langle X \rangle \in \text{DL}(\Lambda_{\mathbb{Z}_{(p)}})$ , then

$$\langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle \wedge \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle X \wedge X \wedge h^\bullet \Lambda_{\mathbb{Q}} \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle,$$

because  $h^\bullet$  maps  $\text{DL}(\Lambda_{\mathbb{Q}})$  into  $\text{DL}(\Lambda_{\mathbb{Z}_{(p)}})$ .

If we consider  $\langle Y \rangle \in \text{DL}(\Lambda_{\mathbb{F}_p})$  and  $\langle Z \rangle \in \text{DL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))$ , then  $\langle g^\bullet Y \rangle$  and  $\langle i_h Z \rangle$  are both in  $\text{DL}(\Lambda_{\mathbb{Z}_{(p)}})$ , so their join is as well.

Now consider the Boolean algebra. Recall that the maximum Bousfield class of  $\text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))$  is  $\langle h^\bullet \Lambda_{\mathbb{Q}} \rangle$ , and it is to this that we require complements to join. Taking  $\langle X \rangle \in \text{BA}(\Lambda_{\mathbb{Z}_{(p)}})$ , with complement  $\langle X^c \rangle$ , we compute

$$\langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle \wedge \langle X^c \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle 0 \rangle, \text{ and}$$

$$\langle X \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle \vee \langle X^c \wedge h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle \Lambda_{\mathbb{Z}_{(p)}} \rangle \wedge \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle = \langle h^\bullet \Lambda_{\mathbb{Q}} \rangle.$$

If we take  $\langle Y \rangle \in \text{BA}(\Lambda_{\mathbb{F}_p})$  then Proposition 4.15 implies  $\langle g^\bullet Y \rangle \in \text{BA}(\Lambda_{\mathbb{Z}_{(p)}})$ . For  $\langle Z \rangle \in \text{BA}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))$ , with complement  $\langle Z^c \rangle$  there, one can show that  $\langle i_h Z \rangle$  is complemented in  $\text{BL}(\Lambda_{\mathbb{Z}_{(p)}})$ , with complement  $\langle i_h(Z^c) \rangle \vee \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle$ . Therefore  $\langle g^\bullet Y \rangle \vee \langle i_h Z \rangle \in \text{BA}(\Lambda_{\mathbb{Z}_{(p)}})$ .  $\square$

**Corollary 5.19.** *The Bousfield lattice of  $D(\Lambda_{\mathbb{Z}_{(p)}})$  has cardinality  $2^{2^{\aleph_0}}$ .*

**Proof.** Corollary B in [DP08] states that the Bousfield lattice of  $D(\Lambda_{\mathbb{F}_p})$  has cardinality  $2^{2^{\aleph_0}}$ , so  $\text{BL}(\Lambda_{\mathbb{Z}_{(p)}})$  is at least as large. However,  $\Lambda_{\mathbb{Z}_{(p)}}$  is countable, so [DP01, Thm. 1.2] implies that  $\text{BL}(\Lambda_{\mathbb{Z}_{(p)}})$  has cardinality at most  $2^{2^{\aleph_0}}$ .  $\square$

Corollary 5.16 shows that  $\text{BL}(\Lambda_{\mathbb{F}_p})$  and  $\text{BL}(\text{loc}(g^\bullet \Lambda_{\mathbb{F}_p}))$  are isomorphic as lattices. However, the relationship between  $\text{BL}(\Lambda_{\mathbb{Q}})$  and  $\text{BL}(\text{loc}(h^\bullet \Lambda_{\mathbb{Q}}))$  appears to be more subtle. We have the following negative result, which suggests that the analog of Theorem 5.9 should fail.

**Lemma 5.20.** *Consider the functors  $h_\bullet : D(\Lambda_{\mathbb{Z}_{(p)}}) \rightleftarrows D(\Lambda_{\mathbb{Q}}) : h^\bullet$ . It is not the case that  $\langle h_\bullet h^\bullet X \rangle = \langle X \rangle$  for all  $\langle X \rangle$ .*

**Proof.** Consider again the  $\Lambda_{\mathbb{Q}}$ -module  $\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}$ , now as an object of  $D(\Lambda_{\mathbb{Q}})$  concentrated in degree zero. Lemma 5.7 shows that we can identify  $\Sigma F$  with  $h^\bullet(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}})$ . This implies

$$\langle h^\bullet(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \rangle = \langle F \rangle = \langle g^\bullet \Lambda_{\mathbb{F}_p} \rangle.$$

In particular,  $h^\bullet(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \wedge h^\bullet \Lambda_{\mathbb{Q}} = 0$ . On the other hand,

$$h^\bullet((\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \wedge \Lambda_{\mathbb{Q}}) = h^\bullet(\Lambda_{\mathbb{Q}}/\Lambda_{\mathbb{Z}_{(p)}}) \neq 0.$$

This shows that the equivalent conditions of Lemma 4.13 fail to hold.  $\square$

## REFERENCES

- [Bir79] Garrett Birkhoff, *Lattice theory*, 3rd ed., American Mathematical Society Colloquium Publications, vol. 25, American Mathematical Society, Providence, R.I., 1979.
- [Bou79a] A. K. Bousfield, *The Boolean algebra of spectra*, Comment. Math. Helv. **54** (1979), no. 3, 368–377.
- [Bou79b] ———, *The localization of spectra with respect to homology*, Topology **18** (1979), no. 4, 257–281.
- [BCR97] D. J. Benson, Jon F. Carlson, and Jeremy Rickard, *Thick subcategories of the stable module category*, Fund. Math. **153** (1997), no. 1, 59–80.

- [BIK08] Dave Benson, Srikanth B. Iyengar, and Henning Krause, *Local cohomology and support for triangulated categories*, Ann. Sci. Éc. Norm. Supér. (4) **41** (2008), no. 4, 573–619.
- [BIK11] ———, *Stratifying triangulated categories*, J. Topol. **4** (2011), no. 3, 641–666.
- [DP01] William G. Dwyer and John H. Palmieri, *Ohkawa’s theorem: there is a set of Bousfield classes*, Proc. Amer. Math. Soc. **129** (2001), no. 3, 881–886.
- [DP08] W. G. Dwyer and J. H. Palmieri, *The Bousfield lattice for truncated polynomial algebras*, Homology Homotopy Appl. **10** (2008), no. 1, 413–436.
- [DS95] W. G. Dwyer and J. Spaliński, *Homotopy theories and model categories*, North-Holland, Amsterdam, 1995, pp. 73–126.
- [Hov99] Mark Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, 1999.
- [HPS97] Mark Hovey, John H. Palmieri, and Neil P. Strickland, *Axiomatic stable homotopy theory*, Mem. Amer. Math. Soc. **128** (1997), no. 610.
- [HP99] Mark Hovey and John H. Palmieri, *The structure of the Bousfield lattice*, Homotopy invariant algebraic structures (Baltimore, MD, 1998), Contemp. Math., vol. 239, Amer. Math. Soc., Providence, RI, 1999, pp. 175–196.
- [IK11] Srikanth B. Iyengar and Henning Krause, *The Bousfield lattice of a triangulated category and stratification* (2011), available at [arXiv:1105.1799v2\[math.RT\]](https://arxiv.org/abs/1105.1799v2). preprint.
- [Kra10] Henning Krause, *Localization theory for triangulated categories*, London Math. Soc. Lecture Note Ser., vol. 375, Cambridge Univ. Press, Cambridge, 2010, pp. 161–235.
- [Nee92] Amnon Neeman, *The chromatic tower for  $D(R)$* , Topology **31** (1992), no. 3, 519–532. With an appendix by Marcel Bökstedt.
- [Nee00] ———, *Oddball Bousfield classes*, Topology **39** (2000), no. 5, 931–935.
- [Nee01] ———, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- [Rav84] Douglas C. Ravenel, *Localization with respect to certain periodic homology theories*, Amer. J. Math. **106** (1984), no. 2, 351–414.
- [Ste12] Greg Stevenson, *Derived categories of absolutely flat rings* (2012), available at [arXiv:1210.0399\[math.AC\]](https://arxiv.org/abs/1210.0399). preprint.
- [Wei94] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.

*E-mail address:* [luke.wolcott@gmail.com](mailto:luke.wolcott@gmail.com)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO N6A5B7, CANADA